

482 INHALATIONAL LUNG DISEASE

220. Schwartz DA, Fuortes LJ, Galvin JR, et al. Asbestos-induced pleural fibrosis and impaired lung function. *Am Rev Respir Dis* 1990;141:321-326.
221. Geffer WB, Conant EF. Issues and controversies in the plain-film diagnosis of asbestos-related disorders in the chest. *J Thorac Imaging* 1988;3:11-28.
222. Anton HC. Multiple pleural plaques. *Br J Radiol* 1967;40:685-690.
223. Wain S, Roggli V, Foster W. Parietal pleural plaques, asbestos bodies, and neoplasia: A clinical, pathologic, and roentgenographic correlation of 25 consecutive cases. *Chest* 1984; 86:707-713.
224. Rudd RM. New developments in asbestos-related pleural disease. *Thorax* 1996;51:210-216.
225. Kannerstein M. Recent advances and perspectives relevant to the pathology of asbestos-related diseases in man. *IARC Sci Publ* 1980;149-162.
226. Sebastien P, Janson X, Gaudichet A, et al. Asbestos retention in human respiratory tissues: comparative measurements in lung parenchyma and in parietal pleura. *IARC Sci Publ* 1980;237-246.
227. Albelda SM, Epstein DM, Geffer WB, et al. Pleural thickening: its significance and relationship to asbestos dust exposure. *Am Rev Respir Dis* 1982;126:621-624.
228. Hilt B, Lien JT, Lund-Larsen PG, et al. Asbestos-related findings in chest radiographs of the male population of the county of Telemark, Norway: a cross-sectional study. *Scand J Work Environ Health* 1986;12:567-573.
229. Sider L, Holland EA, Davis TM Jr, et al. Changes on radiographs of wives of workers exposed to asbestos. *Radiology* 1987;164:723-726.
230. Churg A. Asbestos fibers and pleural plaques in a general autopsy population. *Am J Pathol* 1982;109:88-96.
231. Mollo F, Andron A, Bellis D, et al. Screening of autopsy populations for previous occupational exposure to asbestos. *Arch Environ Health* 1987;42:44-50.
232. Rous V, Studeny J. Aetiology of pleural plaques. *Thorax* 1970;25: 270-284.
233. Baris I, Simonato L, Artvinli M, et al. Epidemiological and environmental evidence of the health effects of exposure to erionite fibres: a four-year study in the Cappadocian region of Turkey. *Int J Cancer* 1987;39:10-17.
234. Stephens M, Gibbs AR, Pooley FD, et al. Asbestos induced diffuse pleural fibrosis: pathology and mineralogy. *Thorax* 1987;42:583-588.
235. Hu H, Beckett L, Kelsey K, et al. The left-sided predominance of asbestos-related pleural disease. *Am Rev Respir Dis* 1993;148:981-984.
236. Gallego JC. Absence of left-sided predominance in asbestos-related pleural plaques: a CT study. *Chest* 1988;113:1034-1036.
237. Rockoff S, Kagan E, Schwartz A, et al. Visceral pleural thickening in asbestos exposure: The occurrence and implications of thickened interlobar fissures. *J Thorac Imag* 1987;2:58-66.
238. Webb WR, Cooper C, Gamsu G. Interlobar pleural plaque mimicking a lung nodule in a patient with asbestos exposure. *J Comput Assist Tomogr* 1983;7:135-136.
239. Lynch D, Gamsu G, Aberle D. Conventional and high resolution CT in the diagnosis of asbestos-related diseases. *RadioGraphics* 1989;9:523-551.
240. Davies D, Andrews MI, Jones JS. Asbestos induced pericardial effusion and constrictive pericarditis. *Thorax* 1991;46:429-432.
241. Cooper M, Johnson K, Delany DJ. Case report: asbestos related pericardial disease. *Clin Radiol* 1996;51:656-657.
242. Yazicioglu S. Pleural calcification associated with exposure to chrysotile asbestos in southeast Turkey. *Chest* 1976;70:43-47.
243. Fischbein L, Namade M, Sachs RN, et al. Chronic constrictive pericarditis associated with asbestosis. *Chest* 1988;94:646-647.
244. Baker E, Greene R. Incremental value of oblique chest radiographs in the diagnosis of asbestos-induced pleural disease. *Am J Ind Med* 1982;3:17-22.
245. Sargent E, Boswell W, Ralls P, et al. Subpleural fat pads in patients exposed to asbestos: Distinction from non-calcified pleural plaques. *Radiology* 1984;152:273-277.
246. Vix VA. Extrapleural costal fat. *Radiology* 1974;112:563-565.
247. Kouris SP, Parker DL, Bender AP, et al. Effects of asbestos-related pleural disease on pulmonary function. *Scand J Work Environ Health* 1991;17:179-183.
248. Broderick A, Fuortes LJ, Merchant JA, et al. Pleural determinants of restrictive lung function and respiratory symptoms in an asbestos-exposed population. *Chest* 1992;101:684-691.
249. Kilburn KH, Warshaw R. Pulmonary functional impairment associated with pleural asbestos disease. Circumscribed and diffuse thickening. *Chest* 1990;98: 965-672.
250. Bourbeau J, Ernst P, Chrome J, et al. The relationship between respiratory impairment and asbestos-related pleural abnormality in an active work force. *Am Rev Respir Dis* 1990;142:837-842.
251. Schwartz DA, Galvin JR, Dayton CS, et al. Determinants of restrictive lung function in asbestos-induced pleural fibrosis. *J Appl Physiol* 1990;68:1932-1937.
252. Shih JF, Wilson JS, Broderick A, et al. Asbestos-induced pleural fibrosis and impaired exercise physiology. *Chest* 1994;105:1370-1376.
253. Aberle DR, Gamsu G, Ray CS, et al. Asbestos-related pleural and parenchymal fibrosis: detection with high-resolution CT. *Radiology* 1988;166:729-734.
254. Friedman AC, Fiel SB, Fisher MS, et al. Asbestos-related pleural disease and asbestosis: A comparison of CT and chest radiography. *AJR Am J Roentgenol* 1988;150:269-275.
255. Kreel L. Computer tomography in the evaluation of pulmonary asbestosis. *Acta Radiol* 1976;17:405-412.
256. Aberle DR, Gamsu G, Ray CS. High-resolution CT of benign asbestos-related diseases: clinical and radiographic correlation. *AJR Am J Roentgenol* 1988;151:883-891.
257. Gevenois PA, de Vuyst P, Dedeire S, et al. Conventional and high-resolution CT in asymptomatic asbestos-exposed workers. *Acta Radiologica* 1994;35: 226-229.
258. Sperber M, Mohan KK. Computed tomography: a reliable diagnostic modality in pulmonary asbestosis. *Comput Radiol* 1984;8:125-132.
259. Copley SJ, Wells AU, Rubens MB, et al. Functional consequences of pleural disease evaluated with chest radiography and CT. *Radiology* 2001;220:237-243.
260. Solomon A. Radiological features of asbestos-related visceral pleural changes. *Am J Ind Med* 1991; 19: 339-355.
261. Solomon A, Irwig L, Sluis-Cremer G, et al. Thickening of pulmonary interlobar fissures: exposure-response relationship in crocidolite and amosite miners. *Br J Ind Med* 1979;36:195-198.
262. Kee ST, Gamsu G, Blanc P. Causes of pulmonary impairment in asbestos-exposed individuals with diffuse pleural thickening. *Am J Respir Crit Care Med* 1996;154:789-793.
263. Blesovsky A. The folded lung. *Br J Dis Chest* 1966;60.
264. Munden RF, Libshitz HI. Rounded atelectasis and mesothelioma. *AJR Am J Roentgenol* 1998;170:1519-1522.
265. Hanke R, Kretzschmar R. Rounded atelectasis. *Semin Roentgenol* 1980;15:174-182.
266. Hillerdal G, Hemmingsson A. Pulmonary pseudotumours and asbestos. *Acta Radiologica* 1979;21:615-620.
267. Menzies R, Fraser R. Round atelectasis. Pathologic and pathogenetic features. *Am J Surg Pathol* 1987;11:674-681.
268. Schneider HJ, Felson B, Gonzales LL. Rounded atelectasis. *AJR Am J Roentgenol* 1980;134:225-232.

269. Voisin C, Fisekci F, Voisin-Saltiel S, et al. Asbestos-related rounded atelectasis. Radiologic and mineralogic data in 23 cases. *Chest* 1995;107:477-481.
270. Carvalho PM, Carr DH. Computed tomography of folded lung. *Clin Radiol* 1990;41:86-91.
271. Doyle TC, Lawler GA. CT features of rounded atelectasis of the lung. *AJR Am J Roentgenol* 1984;143:225-228.
272. Lynch D, Gamsu G, Ray C, et al. Asbestos-related focal lung masses: manifestations on conventional and high-resolution CT scans. *Radiology* 1988;169:603-607.
273. McHugh K, Blaquiére RM. CT features of rounded atelectasis. *AJR Am J Roentgenol* 1989;153:257-260.
274. Taylor PM. Dynamic contrast enhancement of asbestos-related pulmonary pseudotumours. *Br J Radiol* 1988;61:1070-1072.
275. Verschakelen JA, Demaerel P, Coolen J, et al. Rounded atelectasis of the lung: MR appearance. *AJR Am J Roentgenol* 1989;152:965-966.
276. Marchbank ND, Wilson AG, Joseph AE. Ultrasound features of folded lung. *Clin Radiol* 1996;51:433-437.
277. O'Donovan PB, Schenk M, Lim K, et al. Evaluation of the reliability of computed tomographic criteria used in the diagnosis of round atelectasis. *J Thorac Imaging* 1997;12:54-58.
278. Cooke WE. Pulmonary asbestosis. *Br Med J* 1927;2:1024-1025.
279. Walker AM, Loughlin JE, Friedlander ER, et al. Projections of asbestos-related disease 1980-2009. *J Occup Med* 1983; 25:409-425.
280. Craighead JE, Mossman BT. The pathogenesis of asbestos-associated diseases. *N Engl J Med* 1982;306: 1446-1455.
281. Gregor A, Parkes RW, du Bois R, et al. Radiographic progression of asbestosis: preliminary report. *Ann N Y Acad Sci* 1979;330:147-156.
282. Rubino GF, Newhouse M, Murray R, et al. Radiologic changes after cessation of exposure among chrysotile asbestos miners in Italy. *Ann N Y Acad Sci* 1979;330:157-161.
283. Shepherd JR, Hillerdal G, McLarty J. Progression of pleural and parenchymal disease on chest radiographs of workers exposed to amosite asbestos. *Occup Environ Med* 1997;54:410-415.
284. Geffer W, Epstein D, Miller W. Radiographic evaluation of asbestos-related chest disorders. *CRC Crit Rev Diagn Imaging* 1984;21:123-181.
285. Green RA, Dimcheff DG. Massive bilateral upper lobe fibrosis secondary to asbestos exposure. *Chest* 1974;65:52-55.
286. Hillerdal G. Asbestos exposure and upper lobe involvement. *AJR Am J Roentgenol* 1982;139:1163-1166.
287. Feigin DS. Talc: understanding its manifestations in the chest. *AJR Am J Roentgenol* 1986;146:295-301.
288. Staples CA, Gamsu G, Ray CS, et al. High resolution computed tomography and lung function in asbestos-exposed workers with normal chest radiographs. *Am Rev Respir Dis* 1989;139: 1502-1508.
289. Bergin CJ, Castellino RA, Blank N, et al. Specificity of high-resolution CT findings in pulmonary asbestosis: do patients scanned for other indications have similar findings? *AJR Am J Roentgenol* 1994;163:551-555.
290. al Jarad N, Strickland B, Pearson MC, et al. High resolution computed tomographic assessment of asbestosis and cryptogenic fibrosing alveolitis: a comparative study. *Thorax* 1992;47: 645-650.
291. Akira M, Yamamoto S, Inoue Y, et al. High-resolution CT of asbestosis and idiopathic pulmonary fibrosis. *AJR Am J Roentgenol* 2003;181:163-169.
292. Copley S, Wells A, Sivakumaran P, et al. Asbestosis and idiopathic pulmonary fibrosis: comparison thin-section CT features. *Radiology* 2003;229:731-736.
293. Rockoff SD, Schwartz A. Roentgenographic underestimation of early asbestosis by international labor organization classification. *Chest* 1988;93:1988-1991.
294. Oksa P, Suoranta H, Koskinen H, et al. High-resolution computed tomography in the early detection of asbestosis. *Int Arch Occ Envir Health* 1994;65: 299-304.
295. Neri S, Boraschi P, Antonelli A, et al. Pulmonary function, smoking habits, and high resolution computed tomography (HRCT) early abnormalities of lung and pleural fibrosis in shipyard workers exposed to asbestos. *Am J Ind Med* 1996;30:588-595.
296. Akira M, Yokoyama K, Yamamoto S, et al. Early asbestosis: evaluation with high-resolution CT. *Radiology* 1991; 178:409-416.
297. Gamsu G, Salmon CJ, Warnock ML, et al. CT quantification of interstitial fibrosis in patients with asbestosis: a comparison of two methods. *AJR Am J Roentgenol* 1995;164:63-68.
298. Hughes JM, Weill H. Asbestosis as a precursor of asbestos related lung cancer: results of a prospective mortality study. *Br J Ind Med* 1991;48:229-233.
299. Weiss W. Asbestosis: a marker for the increased risk of lung cancer among workers exposed to asbestos. *Chest* 1999;115:536-549.
300. Wilkinson P, Hansell DM, Janssens J, et al. Is lung cancer associated with asbestos exposure when there are no small opacities on the chest radiograph? [see comments]. *Lancet* 1995;345: 1074-1078.
301. Finkelstein MM. Radiographic asbestosis is not a prerequisite for asbestos-associated lung cancer in Ontario asbestos-cement workers. *Am J Ind Med* 1997;32:341-348.
302. Scancarello G, Romeo R, Sartorelli E. Respiratory disease as a result of talc inhalation. *J Occup Environ Med* 1996;38:610-614.
303. Wells IP, Bhatt RC, Flanagan M. Kaolinosis: a radiological review. *Clin Radiol* 1985;36:579-582.
304. Kennedy T, Rawlings W Jr, Baser M, et al. Pneumoconiosis in Georgia kaolin workers. *Am Rev Respir Dis* 1983;127:215-220.
305. Oldham PD. Pneumoconiosis in Cornish china clay workers. *Br J Ind Med* 1983;40:131-137.
306. Han D, Goo JM, Im JG, et al. Thin-section CT findings of arc-welders' pneumoconiosis. *Korean J Radiol* 2000;1:79-83.
307. Akira M. Uncommon pneumoconioses: CT and pathologic findings. *Radiology* 1995;197:403-409.
308. Yoshii C, Matsuyama T, Takazawa A, et al. Welder's pneumoconiosis: diagnostic usefulness of high-resolution computed tomography and ferritin determinations in bronchoalveolar lavage fluid. *Intern Med* 2002;41:1111-1117.
309. Mapel D, Coultas DB. Disorders due to metals other than silica, coal, asbestos, to metals. In: Hendrick D, Burge P, Beckett W, Churg A, eds. *Occupational disorders of the lungs: recognition, management, and prevention*. London: Harcourt, 2002:163-191.
310. Kim CK, Kim SW, Kim JS, et al. Bronchiolitis obliterans in the 1990s in Korea and the United States. *Chest* 2001;120:1101-1106.
311. Shaver CG, Riddell AR. Lung changes associated with the manufacture of alumina abrasive. *J Ind Hyg* 1947;29: 145-157.
312. Brooks SM. Lung disorders resulting from the inhalation of metals. *Clin Chest Med* 1981;2:235-254.
313. Vahlensieck M, Overlack A, Müller KM. Computed tomographic high-attenuation mediastinal lymph nodes after aluminum exposition. *Eur Radiol* 2000;10:1945-1946.
314. Hooper WF. Acute beryllium lung disease. *N C Med J* 1981;42:551-553.
315. Kreiss K, Wasserman S, Mroz MM, et al. Beryllium disease screening in the ceramics industry. Blood lymphocyte test performance and exposure-disease relations. *J Occup Med* 1993;35:267-274.
316. Kreiss K, Mroz MM, Zhen B, et al. Epidemiology of beryllium sensitization and disease in nuclear workers. *Am Rev Respir Dis* 1993;148:985-991.

Unusual causes, including paragonimiasis,⁴²⁷ are described.³⁵⁰ The clinical context in which the effusion occurs is so characteristic that pseudochylothorax rarely causes diagnostic confusion with a true chylothorax. Imaging may also help differentiate between these entities, since true chylothorax is rarely associated with pleural thickening, loculation, or calcification.

HEMOTHORAX

Hemothorax usually results from trauma.⁴²⁸ However, on occasion it occurs in other conditions (Box 15.9). The natural history of hemothorax depends in part on the source of the bleeding. Low pressure bleeding from the lung tends to stop spontaneously because the pleural fluid compresses and collapses the lung. High pressure bleeding from systemic vessels is less susceptible to the tamponade effect of pleural fluid,⁴⁴⁸ and the bleeding may be rapid and persistent with the formation of a tension hemothorax.⁴²⁹ In the context of trauma

Box 15.9 Causes of hemothorax

Trauma

Open

Closed (with or without fracture)

Iatrogenic⁴³¹

Infection

Varicella⁴³²

Coagulopathy

Hemophilia⁴³³

Anticoagulants^{429,434,435}

Vascular abnormality

Arteriovenous malformation⁴³⁶

Dissecting aortic aneurysm

Atherosclerotic aneurysm⁴³⁷

Rib exostosis⁴³⁸

Neurofibromatosis with pregnancy¹³⁰

Pulmonary and pleural neoplasms^{437,439}

Extramedullary hemopoiesis^{440,441}

Pneumothorax^{430,442}

Catamenial hemothorax (endometriosis)^{443,444}

Idiopathic⁴⁴⁵⁻⁴⁴⁷

other causes of rapidly accumulating pleural fluid should be considered, including ruptured esophagus, ruptured thoracic duct, traumatic subarachnoid pleural fistula,⁴⁴⁹ and iatrogenic causes, particularly venous perforation from line placement.

In the acute state nothing on the chest radiograph distinguishes hemothorax from other collections of pleural fluid. However, on CT a hemothorax may show areas of hyperdensity (see Fig. 15.14).⁴⁵⁰ With clotting of the blood, loculation tends to occur and fibrin bodies may form.^{59,430} Hemothorax may eventually organize and cause massive pleural thickening (fibrothorax), necessitating decortication, a

complication that can be avoided by early evacuation of the pleural space.

PLEURAL THICKENING

Pleural thickening can be localized or diffuse, and usually represents the organized end stage of a variety of active processes, particularly infective and noninfective inflammation, hemothorax, and asbestos- and drug-related disease (see Box 15.10). It is virtually always present after thoracotomy and pleurodesis,⁴⁵¹ and may follow irradiation. The most common causes are probably hemothorax, bacterial infection, and tuberculosis.⁵ Particularly marked pleural thickening is seen following tuberculosis or irradiation. Identification of diffuse pleural thickening is important because it is commonly associated with significant restrictive physiologic impairment. It can be important to try to distinguish between diffuse pleural thickening, which may be due to a large variety of causes, and localized pleural plaques which are usually related to asbestos exposure (see p. 1043). On the chest radiograph, diffuse postinflammatory pleural thickening almost always involves the costophrenic sulci (Fig. 15.44), while pleural plaques present with localized areas of soft tissue density along the chest wall. The presence of bilateral abnormality favors asbestos related disease.¹⁰

Box 15.10 Causes of pleural thickening

Infection

Tuberculosis

Empyema

Hemothorax

Asbestos exposure

Surgery or pleurodesis

Radiation

Malignancy

Metastasis

Mesothelioma

Leukemia

Lymphoma

The radiographic changes of diffuse pleural thickening are more commonly unilateral and consist of soft tissue shadowing, characteristically in the more dependent lateral and posterior parts of the chest. There may be radiographic signs of ipsilateral rib enlargement in patients with chronic (particularly tuberculous) pleural disease.⁴⁵² Blunting of the costophrenic angle is common and is often angular, distinguishing it from the more smoothly curvilinear pleural fluid. Decubitus radiographs and ultrasonography are particularly helpful in making this distinction. En face, extensive pleural thickening gives a veil-like opacity that has no clear margins and crosses known pulmonary boundaries. Tangentially, it appears as a soft tissue density immediately inside and parallel to the chest wall, sharply margined on its inner aspect and fading into the soft tissues of the chest wall laterally. Such pleural thickening can extend into and thicken fissures.





Fig. 15.44 Diffuse pleural thickening related to previous pneumonia.

A, Frontal chest radiograph shows smooth pleural thickening extending along the right lateral chest wall, with blunting of the costophrenic sulcus, and marked inferior displacement of the right hilum. **B**, Lateral radiograph shows a posterior mass with vessels curving into it. **C**, CT shows dense pleural thickening, with typical features of round atelectasis, with bronchi and vessels curving into the medial and lateral aspects of the mass, and fissural displacement indicating marked right lower lobe volume loss.

On ultrasound, pleural thickening produces a homogeneously echo dense layer subjacent to the chest wall, but it cannot be reliably detected unless it is about 1 cm or more thick.⁸⁶ There is no posterior echo enhancement, but this is often difficult to assess because the soft tissue-lung interface is normally so reflective.

On CT, pleural thickening is detected as a layer of soft tissue opacity lying at the chest wall-lung interface. It can be detected almost as well with conventional CT as with HRCT, though the latter is more sensitive in assessing asbestos related plaques.⁴⁵³ In addition, HRCT may sometimes clarify equivocal findings on conventional CT.⁴⁵⁴ On HRCT, pleural thickening is best assessed inside ribs where there should be no discernible soft tissue; exceptions to this "rule" are discussed on page 40.⁴⁵⁵ Paravertebrally any thickening of the normally insignificant pleural line is abnormal. HRCT is very sensitive and can detect thickening on the order of 1–2 mm. The extrapleural fat layer, which is normally absent or relatively thin, thickens in chronic pleural disease, particularly with chronic empyema,^{455–458} making appreciation of pleural thickening easier. When this fat has higher density than usual, it suggests that there is active inflammation in the pleural space.⁴⁵⁹ Both the distribution and

morphology⁸⁹ of diffuse pleural thickening are helpful in identifying a cause. In one study the specificity of various CT signs in differentiating a malignant from a benign pleural process was evaluated. The four most useful signs of malignancy (with specificities) were circumferential thickening (100%), nodularity (94%), parietal thickening >1 cm (94%), and mediastinal pleural involvement.⁴⁵⁴ Early experience suggests that positron emission tomography (PET) with ¹⁸F-fluorodeoxyglucose (¹⁸F-FDG) may have a role in the differentiation between malignant and benign pleural disease.⁴⁶⁰ In a study of 23 patients, ¹⁸F-FDG PET correctly identified all 16 cases of malignant pleural infiltration; there was intense uptake of ¹⁸F-FDG in 14 of 16 cases (Fig. 15.45). Seven of nine benign lesions were characterized by an absence of tracer uptake but there was moderate uptake in two patients, one with a parapneumonic effusion and one with tuberculous pleurisy.⁴⁶⁰ Another study found that MRI was more accurate than CT in diagnosis of malignant pleural disease: malignant disease was associated with increased signal intensity on T2-weighted images, and with enhancement following gadolinium administration.⁴⁶¹

Pleural thickening, particularly when generalized, is often correctly dismissed as an inactive residuum. Care, however, must




Fig. 15.45 Activity of PET in malignant pleural disease due to mesothelioma (same patient as in Fig. 15.32). **A**, Chest radiograph shows circumferential pleural thickening. **B**, Coronal MRI confirms the presence of circumferential pleural thickening extending into the costophrenic sulci, with a mass in the major fissure. **C**, Coronal PET image obtained with ^{18}F -fluoro-deoxyglucose shows circumferential increase in metabolic activity, extending into the fissures.

be taken to distinguish it from various active processes, some of which are neoplastic. Although a number of these conditions tend to give plaquelike, nodular, or irregular shadowing (Fig. 15.45), they occasionally closely resemble simple inactive pleural thickening. Disorders to consider include mycetoma related pleural thickening,⁴⁶² mesothelioma, diffuse pleural metastases (e.g. from thymoma⁴⁶³), leukemia,⁴⁶⁴ lymphoma, and Wegener granulomatosis. It should also be remembered that a thick pleural peel occurring following empyema will usually decrease progressively over the subsequent 12 weeks.⁴⁶⁵

There are several uncommon syndromes of idiopathic bilateral pleural thickening. In the entity of cryptogenic bilateral fibrosing pleuritis there is widespread thickening of the pleura preceded by effusions (Fig. 15.46).^{466,467} Patients may have an

elevated erythrocyte sedimentation rate, restrictive lung function, and evidence of rounded atelectasis on CT.⁴⁶⁶ Another uncommon idiopathic syndrome is associated with progressive apical pleural and subpleural fibrosis, with restrictive physiology.⁴⁶⁸ This syndrome may be familial,⁴⁶⁹ and may be associated with renal tubular acidosis.⁴⁷⁰

Mimics

Pleural thickening must be differentiated from apical pleural caps and extrapleural fat. The apical pleural cap, though it looks like pleural thickening, is usually due either to extrapleural fat or to subpleural fibrosis. Older publications refer to serratus



Fig. 15.46 Cryptogenic bilateral fibrosing pleuritis. Prone CT image shows bilateral smooth pleural thickening with bilateral lower lobe volume loss and parenchymal scarring.




Fig. 15.47 Extrapleural fat in a patient treated with steroids. Chest radiograph shows lobulated smooth thickening along the chest wall bilaterally, extending to the lung apices, and extending. Symmetric curvilinear interfaces in the upper chest (arrows) are due to invagination of fat into the major fissures bilaterally.

anterior shadows and rib companion shadows as mimics of pleural plaques on chest radiographs.^{471,472} These are not discussed here because it is very rare for the serratus anterior muscle or ribs to be so sharply outlined by air as to produce a radiographically distinct interface.

Extrapleural fat

Extrapleural fat may generate confusing shadows that can resemble generalized pleural thickening or plaques (Fig. 15.47).^{473,474} The distribution varies from patient to patient. Sometimes the fat is widely distributed, mimicking a pleural peel. At other times, it is localized and develops particularly over the fourth to eighth ribs between the anterior axillary line and the rib angles.⁴⁷³ Excess thoracic fat is more common in obese patients, and associated with mediastinal lipomatosis, but may also be found in thinner individuals. Extrapleural fat may be distinguished from pleural thickening by its symmetry and its undulating outline, often extending to the lung apices but sparing the costophrenic sulci.⁴⁷³ If doubt remains, CT will readily differentiate thickening from fat, but this distinction is not usually clinically important.⁴⁷³




Fig. 15.48 Apical pleural caps. Symmetric soft tissue opacities are projected under both second ribs. They are slightly atypical for pleural caps, being thicker (1 cm) than usual, with some irregularity of their lower margins. The appearance of pleural caps is quite variable.

45 years of age.⁴⁷⁵ The opacity is formed by an apical subpleural scar that is nonspecific and unrelated to tuberculosis.⁴⁷⁸

In contrast to the smooth or undulating outline of the apical cap on the chest radiograph, CT usually shows a subpleural irregular linear abnormality, consistent with the pathologic findings of dense subpleural fibrosis, and sometimes difficult to distinguish from a spiculated lung cancer (Fig. 15.49). Indeed, Yousem⁴⁷⁹ presented a series of 13 such cases which had been resected because of suspected lung cancer.

The differential diagnosis of an idiopathic apical cap includes nongranulomatous and granulomatous (tuberculous, fungal) infection, radiation pleuritis, lymphoma, pleural and extrapleural neoplasms, extrapleural hematoma, prominent subclavian artery, mediastinal lipomatosis,⁴⁸⁰ and apicolateral extrapleural fat.⁴⁸¹ A HRCT study has shown, somewhat surprisingly, that the bulk of the apical "pleural" opacity

Apical pleural cap

An idiopathic apical pleural cap is an irregular, usually homogeneous, soft tissue density that is found at the extreme lung apex (Fig. 15.48).⁴⁷⁵ The lower border is usually sharply margined and may be smoothly curvilinear, tented, or undulating.⁴⁷⁶ Caps are usually <5 mm thick, but the width is variable. In two series caps were about as common unilaterally (11 and 7%) as bilaterally (11 and 12%).^{476,477} When bilateral the caps were usually asymmetric. The frequency of occurrence increases with age: 6.2% up to 45 years of age and 15.9% over




Fig. 15.49 Apical pleural scar. **A**, Chest radiograph shows asymmetric soft tissue thickening at the right apex. **B** and **C**, CT shows irregular subpleural density, with spiculation mimicking lung cancer. The patient was followed for 2 years without evidence of progression.





Fig. 15.50 Pancoast tumor. Chest radiograph shows apical pleural thickening associated with destruction of the posterior left second rib.

associated with previous tuberculosis is caused by a thickened (up to 25 mm) layer of fat between visceral pleura and the endothoracic fascia-innermost intercostal muscle stripe.⁴⁸² This may be related to contraction of the upper lobe, with filling of the extrapleural space by fat. The most important differential diagnosis is with Pancoast tumor, which should be suspected if there is marked asymmetry or nodularity of apical pleural

thickening, if the patient has local pain, and particularly if there is underlying bone destruction (Fig. 15.50).⁴⁸⁰

RADIOLOGIC APPEARANCE FOLLOWING PLEURODESIS

The most common indications for pleurodesis are recurrent pneumothorax, and malignant pleural effusion. The perfect agent for pleurodesis has not yet been found, but talc is increasingly used in place of chemical agents such as tetracycline or bleomycin, because of its higher success rates and lower rate of local and systemic symptoms.^{336,483,484} Talc may be administered either as an aerosolized powder at thoracoscopy,^{485,486} or as a slurry via large- or small-bore chest tubes at the bedside.^{336,487,488}

After pleurodesis, the pleural space usually undergoes a phase of organization, with pleural thickening and loculations evident on chest radiograph; about 60% of patients have radiographically visible pleural thickening at longterm follow up.⁴⁵¹ On CT obtained after talc pleurodesis, the pleural space usually reveals variable degrees of pleural thickening and nodularity, often with a residual loculated effusion. High-attenuation areas representing talc deposits may mimic pleural calcification (Fig. 15.51).⁴⁸⁹ The use of talc as a pleurodesis agent remains controversial, mainly because of recurrent reports of acute respiratory distress syndrome or lung edema occurring after talc pleurodesis.^{485,490-492} Although expert opinion favors its use, the search for a better agent continues.



Fig. 15.51 CT appearances following talc pleurodesis. CT in a patient who had aerosolized talc pleurodesis for a left-sided mesothelioma shows a nodular area of hyperattenuation along the left posterior chest wall. A further nodular collection of talc is seen anterior to the heart.

Box 15.11 Causes of pleural calcification

Infection

Tuberculous empyema

Nontuberculous empyema

Hemothorax

Mineral inhalation

Asbestos (including tremolite talc)

Mica

Zeolites

Miscellaneous

Chronic pancreatitis⁵²⁰

Chronic hemodialysis⁵²¹

Calcified metastasis

Alveolar microlithiasis^{522,523}

PLEURAL CALCIFICATION

Virtually any process that can cause pleural thickening can be responsible for later pleural calcification,⁴⁹³ but in practice calcification is usually due to infection, hemorrhage, or asbestos exposure. The recognized causes are listed in Box 15.11.

Calcification in asbestos inhalation and related conditions is morphologically characteristic,⁴⁹⁴ and is considered on page 451. Calcification following infection and hemorrhage generally cannot be distinguished from each other. Such calcification is usually unilateral and varies from barely detectable to massive (Fig. 15.52). In the latter circumstance it becomes sheetlike and, reflecting the gravitationally determined distribution of the preceding pleural fluid, is often concentrated posterolaterally.⁴⁹³ En face it appears as a hazy veil-like opacity, but in profile it is dense and linear, often parallel to the inner chest wall. The calcification in old empyemas occurs in both visceral and parietal pleura.^{457,495} Sometimes these calcified layers are separated, an observation that can be made on radiographs or more easily on CT (Fig. 15.53).⁴⁵⁷ In a series of 140 calcified fibrothoraces, 15.7% had a persistent effusion that was sandwiched between layers of thickened calcified pleura and was demonstrable on CT by virtue




Fig. 15.52 Pleural calcification. Localized view of right middle and lower zone shows sheetlike calcification. Laterally, where calcification is tangential to the x-ray beam, it is dense and homogeneous, but medially – where it is seen en face – it is more broken up and nodular. There is a 1 cm thick band of noncalcified pleural thickening along the lateral chest wall.

of its attenuation, location, homogeneity, and failure to enhance.⁴⁵⁷ This can be suspected from the plain radiograph with pleural thickening of >2 cm and a double layer of calcification.⁴⁵⁷ Active infection of these loculated collections is manifest by expansion of the pleural opacity and development of an air-fluid level signifying a bronchopleural fistula.⁴⁹⁵




Fig. 15.53 Pleural calcification – old tuberculous empyema. Lenticular pleural opacity shows soft tissue density centrally and is margined by heavy calcification both in the visceral and parietal pleura.

Occasionally postempyema calcification in the pleural space is manifest as a milk of calcium collection. These are often lenticular in shape and surrounded by mildly thickened pleura. On CT they are high density (200–300 HU) and typically homogeneous.⁴⁷⁹

THORACIC SPLENOSIS

Thoracic splenosis occurs when tissue from a traumatized spleen crosses an injured diaphragm and proliferates within the left hemithorax.⁴⁹⁷ It is an uncommon condition⁴⁹⁸; in one prospective review there was evidence of splenic tissue within the thorax in only three of 17 patients who had sustained combined splenic and diaphragmatic injury.⁴⁹⁹ Tissue from a traumatized spleen crosses an injured diaphragm and

proliferates within the left thorax.⁴⁹⁷ The resulting pleural nodules are often multiple and usually <3 cm in diameter, but may be up to 7 cm.⁴⁹⁷ The nodules are implanted on parietal or viscera pleura, including fissures.⁵⁰⁰

On radiologic study, lesions of splenosis usually appear as pleural masses of soft tissue attenuation (Fig. 15.54), but they may appear intraparenchymal both on conventional radiographs and CT.⁴⁹⁷ It is likely that the majority of apparently intrapulmonary lesions have pleural contact, though possibly some have been implanted in a lung laceration rather than on the pleural surface. On CT, the lesion may be lobulated or smooth and of soft tissue density.⁴⁹⁹ On T1- and T2-weighted MRI the masses have been shown to be isointense with paraspinal muscles and subcutaneous fat, respectively.⁴⁹⁹ Should the spleen have been removed at the time of trauma, the absence of Howell-Jolly bodies in a blood film would suggest persisting ectopic splenic activity.

The diagnosis may be confirmed with scintiscans using ^{99m}Tc-sulfur colloid, ^{99m}Tc-labeled heat damaged erythrocytes, or ¹¹¹In-labeled platelets, all of which are taken up by the ectopic splenic tissue.^{497,501–503}

PNEUMOTHORAX

Traditionally, pneumothorax is divided into spontaneous and traumatic types. The most common causes in adults are listed in Box 15.12. Only spontaneous pneumothorax is discussed in this chapter, apart from a brief consideration of pneumothorax associated with mechanical ventilation.

Primary spontaneous pneumothorax

A pneumothorax occurring without an obvious precipitating traumatic event is termed *spontaneous*; if, in addition, the individual is apparently healthy, the pneumothorax is termed





Fig. 15.54 Thoracic splenosis in a patient with a history of previous trauma. **A**, Chest radiograph shows a well-defined soft tissue mass in the left upper chest. **B**, CT confirms a posterior pleural mass. Splenosis was confirmed by sulfur colloid scan.




Fig. 15.71 Central bronchopleural fistula in a 70-year-old woman following left pneumonectomy for a bronchial carcinoma. CT shows a large amount of air in the pneumonectomy space. Air leads from the bronchial stump to the pneumonectomy space (arrow).




Fig. 15.72 Peripheral bronchopleural fistula in a 37-year-old female following wedge resection of the superior segment of the right lower lobe. There is air in the pleural space, but the fistula is not directly demonstrated.

Bronchopleural fistulas complicating infections (Fig. 15.62) are considered in the discussion of empyema (see p. 203). Postsurgical fistulas are considered here. They occur with a frequency of about 2.5–3%^{746,747} and usually develop within 2 weeks of surgery. They should be suspected with the postoperative development of fever, hemoptysis, cough (especially if productive of a large amount of brown sputum), and a persistent large air leak from the pleural drains. Postoperative bronchopleural fistulas are usually associated with infection, and are much more common after surgery performed for pulmonary infections such as tuberculosis.⁷⁴⁸

The chest radiographic signs of bronchopleural fistula following recent pneumonectomy are: (1) a sudden increase in the amount of air in the pneumonectomy space, or in the adjacent chest wall; (2) a decreased amount of fluid; (3) loss of the normal mediastinal shift toward the operated side; and (4) sometimes a contralateral parenchymal opacity due to aspiration of fluid from the pneumonectomy space. Occasionally, unchanged persistence of an airspace following pneumonectomy indicates a fistula. This happens when the residual space is surrounded by pleural fibrosis and scarring so that it cannot change shape.⁷⁴⁶ Extensive scarring may also prevent the mediastinal shift sign from being seen.⁷⁴⁹ It is not uncommon for radiographic signs (increasing pleural air) of bronchopleural

fistula to appear in otherwise well patients who go on without complication or interference to successful obliteration of the pleural cavity.⁷⁵⁰ This is ascribed to a flap valve type of fistula that is self healing. Delayed bronchopleural fistula, occurring after air has been eliminated from the pneumonectomy or lobectomy space, is signaled by the reappearance of air in the pleural space (Fig. 15.71).

A fistula may be detected with ¹³³Xe lung scintigraphy in the washout phase.⁷⁵¹ However, it usually cannot be demonstrated with DTPA aerosol. Injection of water soluble contrast into the relevant bronchus, or into the pleural space may occasionally be helpful. CT can be helpful in depicting the anatomic details of a fistula, particularly in a peripheral bronchopleural fistula which cannot be directly visualized at bronchoscopy.^{741,743} Although CT will directly visualize a peripheral bronchopleural fistula in only 30–50% of cases, it will show a probable cause of the bronchopleural fistula (e.g. peripheral cavity, bulla) in most of the others.^{742,743} Use of thin sections through areas of suspected fistula may be helpful in directly identifying the site of leakage.⁷⁴³ Postoperative peripheral bronchopleural fistulas (Fig. 15.72), or those related to bullae, are less likely to be demonstrated than other types of fistulas.^{742,743}

REFERENCES

1. Zocchi L. Physiology and pathophysiology of pleural fluid turnover. *Eur Respir J* 2002;20:1545–1558.
2. Henschke CI, Davis SD, Romano PM, et al. The pathogenesis, radiologic evaluation, and therapy of pleural effusions. *Radiol Clin North Am* 1989;27:1241–1255.
3. Sahn SA. State of the art. The pleura. *Am Rev Respir Dis* 1988;138:184–234.
4. Agostoni E. Mechanics of the pleural space. *Physiol Rev* 1972;52:57–128.
5. Light RW. *Pleural diseases*, 2nd edn. Philadelphia: Lea and Febiger, 1990.
6. Noppen M, De Waele M, Li R, et al. Volume and cellular content of normal pleural fluid in humans examined by pleural lavage. *Am J Respir Crit Care Med* 2000;162:1023–1106.
7. Moskowitz H, Platt RT, Schachar R, et al. Roentgen visualization of minute pleural effusion. An experimental study to determine the minimum amount of pleural fluid

1078 PLEURA AND PLEURAL DISORDERS

- visible on a radiograph. *Radiology* 1973;109:33-35.
8. Hessen J. Roentgen evaluation of pleural fluid: a study of the localization of free effusions, the potentialities of diagnosing minimal quantities of fluid and its existence under physiologic conditions. *Acta Radiol* 1951;86:7-80.
9. Wiener-Kronish JP, Albertine KH, Licko V, et al. Protein egress and entry rates in pleural fluid and plasma in sheep. *J Appl Physiol* 1984;56:459-463.
10. Müller NL. Imaging of the pleura. *Radiology* 1993;186:297-309.
11. Wang NS. Anatomy and physiology of the pleural space. *Clin Chest Med* 1985;6:3-16.
12. Light RW, Macgregor MI, Luchsinger PC, et al. Pleural effusions: the diagnostic separation of transudates and exudates. *Ann Intern Med* 1972;77:507-513.
13. Romero S, Martinez A, Hernandez L, et al. Light's criteria revisited: consistency and comparison with new proposed alternative criteria for separating pleural transudates from exudates. *Respiration* 2000;67:18-23.
14. Heffner JE, Brown LK, Barbieri CA. Diagnostic value of tests that discriminate between exudative and transudative pleural effusions. Primary Study Investigators. *Chest* 1997;111:970-980.
15. Gazquez I, Porcel JM, Vives M, et al. Comparative analysis of Light's criteria and other biochemical parameters for distinguishing transudates from exudates. *Respir Med* 1998;92:762-765.
16. Heffner JE, Highland K, Brown LK. A meta-analysis derivation of continuous likelihood ratios for diagnosing pleural fluid exudates. *Am J Respir Crit Care Med* 2003;167:1591-1599.
17. Chakko S. Pleural effusion in congestive heart failure. *Chest* 1990;98:521-522.
18. Romero-Candeira S, Fernandez C, Martin C, et al. Influence of diuretics on the concentration of proteins and other components of pleural transudates in patients with heart failure. *Am J Med* 2001;110:681-686.
19. Horowitz ML, Schiff M, Samuels J, et al. Pneumocystis carinii pleural effusion. Pathogenesis and pleural fluid analysis. *Am Rev Respir Dis* 1993;148:232-234.
20. el-Naggar T, Abd-el-Maeboud KH, Abdallah MY, et al. Peripartum pleural effusion. *Respir Med* 1994;88:541-542.
21. Gourgoulanis KI, Karantanis AH, Diminikou G, et al. Benign postpartum pleural effusion. *Eur Respir J* 1995;8:1748-1750.
22. Kumagai-Kurata N, Kunitoh H, Nagamine-Nishizawa M, et al. Idiopathic lobular panniculitis with specific pleural involvement. *Eur Respir J* 1995;8:1613-1615.
23. McNeil KD, Fong KM, Walker QJ, et al. Gorham's syndrome: a usually fatal cause of pleural effusion treated successfully with radiotherapy. *Thorax* 1996;51:1275-1276.
24. Panchal N, Bhagat R, Pant C, et al. Allergic bronchopulmonary aspergillosis: the spectrum of computed tomography appearances. *Respir Med* 1997;91:213-219.
25. Man A, Schwarz Y, Greif J. Pleural effusion as a presenting symptom of ovarian hyperstimulation syndrome. *Eur Respir J* 1997;10:2425-2426.
26. Trudo FJ, Gopez EV, Gupta PK, et al. Pleural effusion due to herpes simplex type II infection in an immunocompromised host. *Am J Respir Crit Care Med* 1997;155:371-373.
27. Bass SN, Ailani RK, Shekar R, et al. Pyogenic vertebral osteomyelitis presenting as exudative pleural effusion: a series of five cases. *Chest* 1998;114:642-647.
28. Jay SJ. Diagnostic procedures for pleural disease. *Clin Chest Med* 1985;6:33-48.
29. Smyrniotis NA, Jederlinic PJ, Irwin RS. Pleural effusion in an asymptomatic patient. Spectrum and frequency of causes and management considerations. *Chest* 1990;97:192-196.
30. Maher GG, Berger HW. Massive pleural effusion: malignant and nonmalignant causes in 46 patients. *Am Rev Respir Dis* 1972;105:458-460.
31. Adams RF, Gleeson FV. Percutaneous image-guided cutting-needle biopsy of the pleura in the presence of a suspected malignant effusion. *Radiology* 2001;219:510-514.
32. Adelman M, Albelda SM, Gottlieb J, et al. Diagnostic utility of pleural fluid eosinophilia. *Am J Med* 1984;77:915-920.
33. Jeanfaivre T, Cimon B, Tolstuchow N, et al. Pleural effusion and toxocariasis. *Thorax* 1996;51:106-107.
34. Ishiura Y, Fujimura M, Nakamura N, et al. Intrapleural corticosteroid injection therapy for post-traumatic eosinophilic pleural effusion. *Respir Med* 1996;90:501-503.
35. Kuhn M, Fitting JW, Leuenberger P. Probability of malignancy in pleural fluid eosinophilia. *Chest* 1989;96:992-994.
36. Hirsch A, Ruffie P, Nebut M, et al. Pleural effusion: laboratory tests in 300 cases. *Thorax* 1979;34:106-112.
37. Storey DD, Dines DE, Coles DT. Pleural effusion. A diagnostic dilemma. *JAMA* 1976;236:2183-2186.
38. Ansari T, Idell S. Management of undiagnosed persistent pleural effusions. *Clin Chest Med* 1998;19:407-417.
39. Ferrer JS, Munoz XG, Orriols RM, et al. Evolution of idiopathic pleural effusion: a prospective, long-term follow-up study. *Chest* 1996;109:1508-1513.
40. Agostoni E, D'Angelo E. Thickness and pressure of the pleural liquid at various heights and with various hydrothoraces. *Respir Physiol* 1969;6:330-342.
41. Gillett D, Ford GT, Anthonisen NR. Shape and regional volume in immersed lung lobes. *J Appl Physiol* 1981;51:1457-1462.
42. Raasch BN, Carsky EW, Lane EJ, et al. Pleural effusion: explanation of some typical appearances. *AJR Am J Roentgenol* 1982;139:899-904.
43. Dandy WE Jr. Incomplete pulmonary interlobar fissure sign. *Radiology* 1978;128:21-25.
44. Heitzman E, editor. Subsegmental anatomy of the lung. In: *The lung: Radiographic-pathologic correlations*. St Louis: CV Mosby, 1984.
45. Oestreich AE, Haley C. Pleural effusion: the thorn sign. Not a rare finding. *Chest* 1981;79:365-366.
46. Pecorari A, Weisbrod GL. Computed tomography of pseudotumoral pleural fluid collections in the azygoesophageal recess. *J Comput Assist Tomogr* 1989;13:803-805.
47. Stark P, Leung A. Effects of lobar atelectasis on the distribution of pleural effusion and pneumothorax. *J Thorac Imaging* 1996;11:145-149.
48. Mintzer RA, Hendrix RW, Johnson CS, et al. The radiologic significance of the left pulmonary ligament. Experience with 26 patients. *Chest* 1979;76:401-405.
49. Rabinowitz JC, Wolf BS. Roentgen significance of the pulmonary ligament. *Radiology* 1966;87:1013-1020.
50. Rigby M, Zylak CJ, Wood LD. The effect of lobar atelectasis on pleural fluid distribution in dogs. *Radiology* 1980;136:603-607.
51. Collins JD, Burwell D, Furmanski S, et al. Minimal detectable pleural effusions. A roentgen pathology model. *Radiology* 1972;105:51-53.
52. Vix VA. Roentgenographic recognition of pleural effusion. *JAMA* 1974;229:695-698.
53. Kocijancic I, Terceij M, Vidmar K, et al. The value of inspiratory-expiratory lateral decubitus views in the diagnosis of small pleural effusions. *Clin Radiol* 1999;54:595-597.
54. Petersen J. Recognition of intrapulmonary pleural effusion. *Radiology* 1960;74:34-41.
55. Fleischner F. Atypical arrangement of free pleural effusion. *Radiol Clin North Am* 1963;1:347-362.
56. Bryk D. Intrapulmonary effusion. Effect of expiration on the pseudodiaphragmatic contour. *Radiology* 1976;120:33-36.
57. Schwarz MI, Marmorstein BL. A new radiologic sign of subpulmonic effusion. *Chest* 1975;67:176-178.
58. Dunbar J, Favreau M. Intrapulmonary pleural effusions with particular

- reference to its occurrence in nephrosis. *J Can Assoc Radiol* 1959;10:24.
59. Felson B. Chest roentgenology. Philadelphia: WB Saunders, 1973.
 60. Kaplan LM, Epstein SK, Schwartz SL, et al. Clinical, echocardiographic, and hemodynamic evidence of cardiac tamponade caused by large pleural effusions. *Am J Respir Crit Care Med* 1995;151:904-908.
 61. Lai CL, Tsai TT, Ko SC, et al. Superior vena cava syndrome caused by encapsulated pleural effusion. *Eur Respir J* 1997;10:1675-1677.
 62. Romero S, Martin C, Hernandez L, et al. Effect of body position on gas exchange in patients with unilateral pleural effusion: influence of effusion volume. *Respir Med* 1995;89:297-301.
 63. Agusti AG, Cardus J, Roca J, et al. Ventilation-perfusion mismatch in patients with pleural effusion: effects of thoracentesis. *Am J Respir Crit Care Med* 1997;156:1205-1209.
 64. Pantoja E, Kattan KR, Thomas HA. Some uncommon lower mediastinal densities: a pictorial essay. *Radiol Clin North Am* 1984;22:633-646.
 65. Proto AV, Merhar GL. Central bronchial displacement with large posterior pleural collections. Findings on the lateral chest radiograph and CT scans. *J Can Assoc Radiol* 1984;35:128-132.
 66. Swingle JD, Logan R, Juhl JH. Inversion of the left hemidiaphragm. *JAMA* 1969;208:863-864.
 67. Abu Yousef MM. Case of the fall season. *Semin Roentgenol* 1980;15:269-271.
 68. Lowe SH, Cosgrove DO, Joseph AE. Inversion of the right hemidiaphragm shown on ultrasound examination. *Br J Radiol* 1981;54:754-757.
 69. Subramanyam BR, Raghavendra BN, Lefleur RS. Sonography of the inverted right hemidiaphragm. *AJR Am J Roentgenol* 1981;136:1004-1006.
 70. Katzen BT, Choi WS, Friedman MH, et al. Pseudomass of the liver due to pleural effusion and inversion of the diaphragm. *AJR Am J Roentgenol* 1978;131:1077-1078.
 71. Hertzanu Y, Solomon A. Inversion of the right diaphragm: a thoracoabdominal CT pitfall. *Gastrointest Radiol* 1986;11:200-202.
 72. Levitt RG, Sagel SS, Stanley RJ, et al. Accuracy of computed tomography of the liver and biliary tract. *Radiology* 1977;124:123-128.
 73. Rogers CI, Meredith HC. Osler revisited: an unusual cause of inversion of the diaphragm. *Radiology* 1977;125:596.
 74. Demos TC, Pieters C. Abdominal pseudotumor due to inverted hemidiaphragm. Radiographic, sonographic, and computed tomographic diagnosis. *Chest* 1984;86:466-468.
 75. Dallemand S, Twersky J, Gordon DH. Pseudomass of the left upper quadrant from inversion of the left hemidiaphragm: CT diagnosis. *Gastrointest Radiol* 1982;7:57-59.
 76. Felson B. The extrapleural space. *Semin Roentgenol* 1977;12:327-333.
 77. Feder B, Wilk S. Localized interlobar effusion in heart failure: phantom lung tumor. *Dis Chest* 1956;30:289-297.
 78. Weiss W, Boucot K, Geffer W. Localized interlobar effusion in congestive heart failure. *Ann Intern Med* 1953;38:1177-1186.
 79. Ruskin JA, Gurney JW, Thorsen MK, et al. Detection of pleural effusions on supine chest radiographs. *AJR Am J Roentgenol* 1987;148:681-683.
 80. Woodring JH. Recognition of pleural effusion on supine radiographs: how much fluid is required? *AJR Am J Roentgenol* 1984;142:59-64.
 81. Moller A. Pleural effusion. Use of the semi-supine position for radiographic detection. *Radiology* 1984;150:245-249.
 82. Rudikoff JC. Early detection of pleural fluid. *Chest* 1980;77:109-111.
 83. Trackler RT, Brinker RA. Widening of the left paravertebral pleural line on supine chest roentgenograms in free pleural effusions. *Am J Roentgenol Radium Ther Nucl Med* 1966;96:1027-1034.
 84. Onik G, Goodman PC, Webb WR, et al. Hydropneumothorax: detection on supine radiographs. *Radiology* 1984;152:31-34.
 85. Kuhlman JE, Singha NK. Complex disease of the pleural space: radiographic and CT evaluation. *Radiographics* 1997;17:63-79.
 86. McLoud TC, Flower CD. Imaging the pleura: sonography, CT, and MR imaging. *AJR Am J Roentgenol* 1991;156:1145-1153.
 87. Pugatch RD, Faling LJ, Robbins AH, et al. Differentiation of pleural and pulmonary lesions using computed tomography. *J Comput Assist Tomogr* 1978;2:601-606.
 88. Bressler EL, Francis IR, Glazer GM, et al. Bolus contrast medium enhancement for distinguishing pleural from parenchymal lung disease: CT features. *J Comput Assist Tomogr* 1987;11:436-440.
 89. Maffessanti M, Tommasi M, Pellegrini P. Computed tomography of free pleural effusions. *Eur J Radiol* 1987;7:87-90.
 90. Kollins SA. Computed tomography of the pulmonary parenchyma and chest wall. *Radiol Clin North Am* 1977;15:297-308.
 91. Vock P, Effmann EL, Hedlund LW, et al. Analysis of the density of pleural fluid analogs by computed tomography. *Invest Radiol* 1984;19:10-15.
 92. Williford ME, Hidalgo H, Putman CE, et al. Computed tomography of pleural disease. *AJR Am J Roentgenol* 1983;140:909-914.
 93. Lawton F, Blackledge G, Johnson R. Co-existent chyloous and serous pleural effusions associated with ovarian cancer: a case report of Contarini's syndrome. *Eur J Surg Oncol* 1985;11:177-178.
 94. Sullivan KL, Steiner RM, Wechsler RJ. Lymphaticopleural fistula: diagnosis by computed tomography. *J Comput Assist Tomogr* 1984;8:1005-1006.
 95. Aquino SL, Webb WR, Gushiken BJ. Pleural exudates and transudates: diagnosis with contrast-enhanced CT. *Radiology* 1994;192:803-808.
 96. Arenas-Jimenez J, Alonso-Charterina S, Sanchez-Paya J, et al. Evaluation of CT findings for diagnosis of pleural effusions. *Eur Radiol* 2000;10:681-690.
 97. Dwyer A. The displaced crus: a sign for distinguishing between pleural fluid and ascites on computed tomography. *J Comput Assist Tomogr* 1978;2:598-599.
 98. Naidich DP, Megibow AJ, Hilton S, et al. Computed tomography of the diaphragm: peridiaphragmatic fluid localization. *J Comput Assist Tomogr* 1983;7:641-649.
 99. Halvorsen RA, Fedyschin PJ, Korobkin M, et al. CT differentiation of pleural effusion from ascites. An evaluation of four signs using blinded analysis of 52 cases. *Invest Radiol* 1986;21:391-395.
 100. Teplick JG, Teplick SK, Goodman L, et al. The interface sign: a computed tomographic sign for distinguishing pleural and intra-abdominal fluid. *Radiology* 1982;144:359-362.
 101. Alexander ES, Proto AV, Clark RA. CT differentiation of subphrenic abscess and pleural effusion. *AJR Am J Roentgenol* 1983;140:47-51.
 102. Federle MP, Mark AS, Guillaumin ES. CT of subpulmonic pleural effusions and atelectasis: criteria for differentiation from subphrenic fluid. *AJR Am J Roentgenol* 1986;146:685-689.
 103. Silverman PM, Baker ME, Mahony BS. Atelectasis and subpulmonic fluid: a CT pitfall in distinguishing pleural from peritoneal fluid. *J Comput Assist Tomogr* 1985;9:763-766.
 104. Griffin DJ, Gross BH, McCracken S, et al. Observations on CT differentiation of pleural and peritoneal fluid. *J Comput Assist Tomogr* 1984;8:24-28.
 105. Stark DD, Federle MP, Goodman PC, et al. Differentiating lung abscess and empyema: radiography and computed tomography. *AJR Am J Roentgenol* 1983;141:163-167.
 106. Hirsch JH, Rogers JV, Mack LA. Real-time sonography of pleural opacities. *AJR Am J Roentgenol* 1981;136:297-301.
 107. Yang PC, Luh KT, Chang DB, et al. Value of sonography in determining the nature of pleural effusion: analysis of 320 cases. *AJR Am J Roentgenol* 1992;159:29-33.
 108. Marks WM, Filly RA, Callen PW. Real-time evaluation of pleural lesions: new

1080 PLEURA AND PLEURAL DISORDERS

- observations regarding the probability of obtaining free fluid. *Radiology* 1982;142:163-164.
109. Rosenberg ER. Ultrasound in the assessment of pleural densities. *Chest* 1983;84:283-285.
 110. Lomas DJ, Padley SP, Flower CD. The sonographic appearances of pleural fluid. *Br J Radiol* 1993;66:619-624.
 111. Wu RG, Yuan A, Liaw YS, et al. Image comparison of real-time gray-scale ultrasound and color Doppler ultrasound for use in diagnosis of minimal pleural effusion. *Am J Respir Crit Care Med* 1994;150:510-514.
 112. Wu RG, Yang PC, Kuo SH, et al. "Fluid color" sign: a useful indicator for discrimination between pleural thickening and pleural effusion. *J Ultrasound Med* 1995;14:767-769.
 113. Lipscomb DJ, Flower CD, Hadfield JW. Ultrasound of the pleura: an assessment of its clinical value. *Clin Radiol* 1981;32:289-290.
 114. Dorne HL. Differentiation of pulmonary parenchymal consolidation from pleural disease using the sonographic fluid bronchogram. *Radiology* 1986;158:41-42.
 115. Yu CJ, Yang PC, Wu HD, et al. Ultrasound study in unilateral hemithorax opacification. Image comparison with computed tomography. *Am Rev Respir Dis* 1993;147:430-434.
 116. O'Moore PV, Mueller PR, Simeone JF, et al. Sonographic guidance in diagnostic and therapeutic interventions in the pleural space. *AJR Am J Roentgenol* 1987;149:1-5.
 117. Weingardt JP, Guico RR, Nemcek AA Jr, et al. Ultrasound findings following failed, clinically directed thoracenteses. *J Clin Ultrasound* 1994;22:419-426.
 118. Grogan DR, Irwin RS, Channick R, et al. Complications associated with thoracentesis. A prospective, randomized study comparing three different methods. *Arch Intern Med* 1990;150:873-877.
 119. Osranek M, Bursi F, O'Leary PW, et al. Hand-carried ultrasound-guided pericardiocentesis and thoracentesis. *J Am Soc Echocardiogr* 2003;16:480-484.
 120. Petersen S, Freitag M, Albert W, et al. Ultrasound-guided thoracentesis in surgical intensive care patients. *Intensive Care Med* 1999;25:1029.
 121. Lichtenstein D, Hulot JS, Rabiller A, et al. Feasibility and safety of ultrasound-aided thoracentesis in mechanically ventilated patients. *Intensive Care Med* 1999;25:955-958.
 122. Keske U. Ultrasound-aided thoracentesis in intensive care patients. *Intensive Care Med* 1999;25:896-897.
 123. Himelman RB, Callen PW. The prognostic value of loculations in parapneumonic pleural effusions. *Chest* 1986;90:852-856.
 124. Newlin N, Silver TM, Stuck KJ, et al. Ultrasonic features of pyogenic liver abscesses. *Radiology* 1981;139:155-159.
 125. Patz EJ, Shaffer K, Piwnica WD, et al. Malignant pleural mesothelioma: value of CT and MR imaging in predicting resectability. *AJR Am J Roentgenol* 1992;159:961-966.
 126. Imanaka K, Sugimoto K, Aoki M, et al. MRI finding of chronic hemorrhagic empyema: a case report. *Radiat Med* 1996;14:201-203.
 127. Tscholakoff D, Sechtem U, de Geer G, et al. Evaluation of pleural and pericardial effusions by magnetic resonance imaging. *Eur J Radiol* 1987;7:169-174.
 128. Joseph J, Sahn SA. Connective tissue diseases and the pleura. *Chest* 1993;104:262-270.
 129. Hughson WG, Friedman PJ, Feigin DS, et al. Postpartum pleural effusion: a common radiologic finding. *Ann Intern Med* 1982;97:856-858.
 130. Heffner JE, Sahn SA. Pleural disease in pregnancy. *Clin Chest Med* 1992;13:667-678.
 131. Haddad B, Barton JR, Livingston JC, et al. HELLP (hemolysis, elevated liver enzymes, and low platelet count) syndrome versus severe preeclampsia: onset at < or = 28.0 weeks' gestation. *Am J Obstet Gynecol* 2000;183:1475-1479.
 132. Sahn SA, Miller KS. Obscure pleural effusion. Look to the kidney. *Chest* 1986;90:631.
 133. Sanders RC. Post-operative pleural effusion and subphrenic abscess. *Clin Radiol* 1970;21:308-312.
 134. DeCrosse JJ, Poulin TL, Fox PS, et al. Subphrenic abscess. *Surg Gynecol Obstet* 1974;138:841-846.
 135. Sherman NJ, Davis JR, Jesseph JE. Subphrenic abscess. A continuing hazard. *Am J Surg* 1969;117:117-123.
 136. Miller WT, Talman EA. Subphrenic abscess. *Am J Roentgenol Radium Ther Nucl Med* 1967;101:961-969.
 137. Ballantyne KC, Sethia B, Reece IJ, et al. Empyema following intra-abdominal sepsis. *Br J Surg* 1984;71:723-725.
 138. Haaga JR, Weinstein AJ. CT-guided percutaneous aspiration and drainage of abscesses. *AJR Am J Roentgenol* 1980;135:1187-1194.
 139. Rubin RH, Swartz MN, Malt R. Hepatic abscess: changes in clinical, bacteriologic and therapeutic aspects. *Am J Med* 1974;57:601-610.
 140. Light RW. Exudative pleural effusions secondary to gastrointestinal diseases. *Clin Chest Med* 1985;6:103-111.
 141. Logue R, Rogers J, Gay B. Subtle radiographic signs of left heart failure. *Am Heart J* 1963;65:464-473.
 142. McPeak E, Levine S. The preponderance of right hydrothorax in congestive heart failure. *Ann Intern Med* 1946;25:916-927.
 143. Leuallen E, Carr D. Pleural effusion: a statistical study of 436 patients. *N Engl J Med* 1955;252:79-83.
 144. Bedford D, Lovibond J. Hydrothorax in heart failure. *Br Heart J* 1941;3:93-111.
 145. Peterman TA, Brothers SK. Pleural effusions in congestive heart failure and in pericardial disease. *N Engl J Med* 1983;309:313.
 146. Weiss JM, Spodick DH. Laterality of pleural effusions in chronic congestive heart failure. *Am J Cardiol* 1984;53:951.
 147. White P, August S, Michie C. Hydrothorax in congestive heart failure. *Am J Med Sci* 1947;214:243-247.
 148. Race G, Scheifley C, Edwards J. Hydrothorax in congestive heart failure. *Am J Med* 1957;22:83-89.
 149. Mellins RB, Levine OR, Fishman AP. Effect of systemic and pulmonary venous hypertension on pleural and pericardial fluid accumulation. *J Appl Physiol* 1970;29:564-569.
 150. Wiener-Kronish JP, Matthay MA, Callen PW, et al. Relationship of pleural effusions to pulmonary hemodynamics in patients with congestive heart failure. *Am Rev Respir Dis* 1985;132:1253-1256.
 151. Wiener-Kronish JP, Goldstein R, Matthay RA, et al. Lack of association of pleural effusion with chronic pulmonary arterial and right atrial hypertension. *Chest* 1987;92:967-970.
 152. Lee YC, Vaz MA, Ely KA, et al. Symptomatic persistent post-coronary artery bypass graft pleural effusions requiring operative treatment: clinical and histologic features. *Chest* 2001;119:795-800.
 153. Light RW. Pleural effusions after coronary artery bypass graft surgery. *Curr Opin Pulm Med* 2002;8:308-311.
 154. Kollef MH. Chronic pleural effusion following coronary artery revascularization with the internal mammary artery. *Chest* 1990;97:750-751.
 155. Parenti C. Pulmonary embolism after coronary artery bypass surgery. *Crit Care Nurs Q* 1994;17:48-50.
 156. Pego-Fernandes PM, Ebaid GX, Nouer GH, et al. Chylothorax after myocardial revascularization with the left internal thoracic artery. *Arq Bras Cardiol* 1999;73:383-390.
 157. Brancaccio G, Prifti E, Cricco AM, et al. Chylothorax: a complication after internal thoracic artery harvesting. *Ital Heart J* 2001;2:559-562.
 158. Kaminsky ME, Rodan BA, Osborne DR, et al. Postpericardiotomy syndrome. *AJR Am J Roentgenol* 1982;138:503-508.
 159. Weiss JM, Spodick DH. Association of left pleural effusion with pericardial disease. *N Engl J Med* 1983;308:696-697.
 160. Dressler W. The post-myocardial-infarction syndrome: a report on 44 cases. *Arch Intern Med* 1959;103:28-42.

161. Engle M, Ito T. The postpericardiotomy syndrome. *Am J Cardiol* 1961;7:73-82.
162. Hearne C, Forjuoh SN. Postcardiac injury syndrome after coronary angioplasty and stenting. *J Am Board Fam Pract* 2003;16:73-74.
163. Stefanelli CB, Bradley DJ, Leroy S, et al. Implantable cardioverter defibrillator therapy for life-threatening arrhythmias in young patients. *J Interv Card Electrophysiol* 2002;6:235-244.
164. Spindler M, Burrows G, Kowalik P, et al. Postpericardiotomy syndrome and cardiac tamponade as a late complication after pacemaker implantation. *Pacing Clin Electrophysiol* 2001;24:1433-1434.
165. Bajaj BP, Evans KE, Thomas P. Postpericardiotomy syndrome following temporary and permanent transvenous pacing. *Postgrad Med J* 1999;75:357-358.
166. Liem KL, ten Veen JH, Lie KI, et al. Incidence and significance of heartmuscle antibodies in patients with acute myocardial infarction and unstable angina. *Acta Med Scand* 1979;206:473-475.
167. Sahn SA. Immunologic diseases of the pleura. *Clin Chest Med* 1985;6:83-102.
168. Stelzner TJ, King TE Jr, Antony VB, et al. The pleuropulmonary manifestations of the postcardiac injury syndrome. *Chest* 1983;84:383-387.
169. Areno JP, McCartney JP, Eggerstedt J, et al. Persistent pleural effusions following coronary bypass surgery. *Chest* 1998;114:311-474.
170. Levin EJ, Bryk D. Dressler syndrome (postmyocardial infarction syndrome). *Radiology* 1966;87:731-736.
171. Tabatznik B, Isaacs J. Postpericardiotomy syndrome following traumatic hemopericardium. *Am J Cardiol* 1961;7:83-96.
172. Soulen RL, Freeman E. Radiologic evaluation of myocardial infarction. *Radiol Clin North Am* 1971;9:567-582.
173. Szabo G, Magyar Z. Effect of increased systemic venous pressure on lymph pressure and flow. *Am J Physiol* 1967;212:1469-1474.
174. Hussey H, Katz S, Yater W. The superior vena caval syndrome: report of thirty-five cases. *Am Heart J* 1946;31:1-26.
175. Perez CA, Presant CA, Van Amburg AL, 3rd. Management of superior vena cava syndrome. *Semin Oncol* 1978;5:123-134.
176. Javaheri S, Hales CA. Sarcoidosis: a cause of innominate vein obstruction and massive pleural effusion. *Lung* 1980;157:81-85.
177. Good JT Jr, Moore JB, Fowler AA, et al. Superior vena cava syndrome as a cause of pleural effusion. *Am Rev Respir Dis* 1982;125:246-247.
178. Diaconis JN, Weiner CI, White DW. Primary subclavian vein thrombosis and bilateral chylothorax documented by lymphography and venography. *Radiology* 1976;119:557-558.
179. Seibert JJ, Golladay ES, Keller C. Chylothorax secondary to superior vena caval obstruction. *Pediatr Radiol* 1982;12:252-254.
180. Bynum LJ, Wilson JE 3rd. Radiographic features of pleural effusions in pulmonary embolism. *Am Rev Respir Dis* 1978;117:829-834.
181. Moses DC, Silver TM, Bookstein JJ. The complementary roles of chest radiography, lung scanning, and selective pulmonary angiography in the diagnosis of pulmonary embolism. *Circulation* 1974;49:179-188.
182. Talbot S, Worthington BS, Roebuck EJ. Radiographic signs of pulmonary embolism and pulmonary infarction. *Thorax* 1973;28:198-203.
183. Dalen JE, Haffajee CI, Alpert JS 3rd, et al. Pulmonary embolism, pulmonary hemorrhage and pulmonary infarction. *N Engl J Med* 1977;296:1431-1435.
184. Simon HB, Daggett WM, DeSanctis RW. Hemothorax as a complication of anticoagulant therapy in the presence of pulmonary infarction. *JAMA* 1969;208:1830-1834.
185. McArdle CR, Sacks BA. Ovarian hyperstimulation syndrome. *AJR Am J Roentgenol* 1980;135:835-836.
186. Schenker JG, Weinstein D. Ovarian hyperstimulation syndrome: a current survey. *Fertil Steril* 1978;30:255-268.
187. Miller WT Jr. Drug-related pleural and mediastinal disorders. *J Thorac Imaging* 1991;6:36-51.
188. Xiol X, Guardiola J. Hepatic hydrothorax. *Curr Opin Pulm Med* 1998;4:239-242.
189. Kinasevitz GT, Keddissi JI. Hepatic hydrothorax. *Curr Opin Pulm Med* 2003;9:261-265.
190. Morrow C, Kantor M, Armen R. Hepatic hydrothorax. *Ann Intern Med* 1958;49:193-203.
191. Islam N, Ali S, Kabir H. Hepatic hydrothorax. *Br J Dis Chest* 1965;59:222-227.
192. Mattison LE, Coppage L, Alderman DF, et al. Pleural effusions in the medical ICU: prevalence, causes, and clinical implications. *Chest* 1997;111:1018-1023.
193. Johnston R, Loo R. Hepatic hydrothorax: studies to determine the source of the fluid and report of thirteen cases. *Ann Intern Med* 1964;61:385-401.
194. Lieberman FL, Hidemura R, Peters RL, et al. Pathogenesis and treatment of hydrothorax complicating cirrhosis with ascites. *Ann Intern Med* 1966;64:341-351.
195. Lieberman FL, Peters RL. Cirrhotic hydrothorax. Further evidence that an acquired diaphragmatic defect is at fault. *Arch Intern Med* 1970;125:114-117.
196. McKay D, Sparling H, Robbins S. Cirrhosis of the liver with massive hydrothorax. *Arch Intern Med* 1947;79:501-509.
197. Williams M. Pleural effusion produced by abdominal-pleural communication in a patient with Laennec's cirrhosis of the liver and ascites. *Ann Intern Med* 1950;33:216-221.
198. Mirouze D, Juttner HU, Reynolds TB. Left pleural effusion in patients with chronic liver disease and ascites. Prospective study of 22 cases. *Dig Dis Sci* 1981;26:984-988.
199. Black LF. The pleural space and pleural fluid. *Mayo Clin Proc* 1972;47:493-506.
200. Frazer IH, Lichtenstein M, Andrews JT. Pleuroperitoneal effusion without ascites. *Med J Aust* 1983;2:520-521.
201. Bradley JW, Feilding LP. Hydropneumothorax complicating perforated peptic ulcer. *Br J Surg* 1972;59:72-73.
202. Nakamura A, Kojima Y, Ohmi H, et al. Peritoneal-pleural communications in hepatic hydrothorax demonstrated by thoracoscopy. *Chest* 1996;109:579-581.
203. Siegerstetter V, Deibert P, Ochs A, et al. Treatment of refractory hepatic hydrothorax with transjugular intrahepatic portosystemic shunt: long-term results in 40 patients. *Eur J Gastroenterol Hepatol* 2001;13:529-534.
204. Spencer EB, Cohen DT, Darcy MD. Safety and efficacy of transjugular intrahepatic portosystemic shunt creation for the treatment of hepatic hydrothorax. *J Vasc Interv Radiol* 2002;13:385-390.
205. Salmon V. Benign pelvic tumours associated with ascites and pleural effusion. *J Mt Sinai Hosp* 1934;1:169-172.
206. Meigs J, Cass J. Fibroma of the ovary with ascites and hydrothorax: with a report of seven cases. *Am J Obstet Gynecol* 1937;33:249-267.
207. Meigs J. Fibroma of the ovary with ascites and hydrothorax Meigs' syndrome. *Am J Obstet Gynecol* 1954;67:962-987.
208. Majzlin C, Stevens F. Meigs' syndrome: case report and review of literature. *J Int Coll Surg* 1964;42:625-630.
209. Solomon S, Farber S, Caruso L. Fibromyomata of the uterus with hemothorax - Meigs' syndrome? *Arch Intern Med* 1971;127:307-309.
210. Kebapci M, Aslan O, Kaya T, et al. Pedunculated uterine leiomyoma associated with pseudo-Meigs' syndrome and elevated CA-125 level: CT features. *Eur Radiol* 2002;12 Suppl 3:S127-129.
211. La Fianza A, Alberici E. CT diagnosis of Pseudo-Meigs' syndrome. *Clin Radiol* 2002;57:315-317.
212. Weise M, Westphalen S, Fayyazi A, et al. Pseudo-Meigs syndrome: uterine

1082 PLEURA AND PLEURAL DISORDERS

- leiomyoma with bladder attachment associated with ascites and hydrothorax – a rare case of a rare syndrome. *Onkologie* 2002;25:443–446.
213. Amant F, Gabriel C, Timmerman D, et al. Pseudo-Meigs' syndrome caused by a hydropic degenerating uterine leiomyoma with elevated CA 125. *Gynecol Oncol* 2001;83:153–157.
 214. Migishima F, Jobo T, Hata H, et al. Uterine leiomyoma causing massive ascites and left pleural effusion with elevated CA 125: a case report. *J Obstet Gynaecol Res* 2000;26:283–287.
 215. Handler CE, Fray RE, Snashall PD. Atypical Meigs' syndrome. *Thorax* 1982;37:396–397.
 216. O'Flanagan SJ, Tighe BF, Egan TJ, et al. Meigs' syndrome and pseudo-Meigs' syndrome. *J R Soc Med* 1987;80:252–253.
 217. Meigs J. Pelvic tumours other than fibromas of the ovary with ascites and hydrothorax. *Obstet Gynecol* 1954;3: 471–485.
 218. Dockerty M, Masson J. Ovarian fibromas: a clinical and pathologic study of 283 cases. *Am J Obstet Gynecol* 1944;47:741–752.
 219. Mokrohisky J. So-called "Meigs' syndrome" associated with benign and malignant ovarian tumors. *Radiology* 1958;70:578–581.
 220. Fishbein R, Murphy G, Wilder R. The pleuropulmonary manifestations of pancreatitis. *Dis Chest* 1962;41:392–397.
 221. Murphy D, Duncan JG, Imrie CW. The "negative chest radiograph" in acute pancreatitis. *Br J Radiol* 1977;50:264–265.
 222. Heller SJ, Noordhoek E, Tenner SM, et al. Pleural effusion as a predictor of severity in acute pancreatitis. *Pancreas* 1997;15: 222–225.
 223. Millward SF, Breatnach E, Simpkins KC, et al. Do plain films of the chest and abdomen have a role in the diagnosis of acute pancreatitis? *Clin Radiol* 1983;34:133–137.
 224. Talamini G, Uomo G, Pezzilli R, et al. Serum creatinine and chest radiographs in the early assessment of acute pancreatitis. *Am J Surg* 1999;177:7–14.
 225. Kaye MD. Pleuropulmonary complications of pancreatitis. *Thorax* 1968;23:297–306.
 226. Rockey DC, Cello JP. Pancreaticopleural fistula. Report of 7 patients and review of the literature. *Medicine (Baltimore)* 1990;69:332–344.
 227. McKenna JM, Chandrasekhar AJ, Skorton D, et al. The pleuropulmonary complications of pancreatitis. Clinical conference in pulmonary disease from Northwestern University-McGaw Medical Center and Veterans Administration Lakeside Hospital, Chicago. *Chest* 1977;71:197–204.
 228. Anderson WJ, Skinner DB, Zuidema GD, et al. Chronic pancreatic pleural effusions. *Surg Gynecol Obstet* 1973;137:827–830.
 229. Miridjanian A, Ambruoso VN, Derby BM, et al. Massive bilateral hemorrhagic pleural effusions in chronic relapsing pancreatitis. *Arch Surg* 1969;98:62–66.
 230. Tewari SC, Jayaswal R, Chauhan MS, et al. Bilateral recurrent haemorrhagic pleural effusion in asymptomatic chronic pancreatitis. *Thorax* 1989;44:824–825.
 231. Cameron JL. Chronic pancreatic ascites and pancreatic pleural effusions. *Gastroenterology* 1978;74:134–140.
 232. Kirchner SG, Heller RM, Smith CW. Pancreatic pseudocyst of the mediastinum. *Radiology* 1977;123:37–42.
 233. Louie S, McGahan JP, Frey C, et al. Pancreatic pleuropericardial effusions. Fistulous tracts demonstrated by computed tomography. *Arch Intern Med* 1985;145:1231–1234.
 234. Faling LJ, Gerzof SG, Daly BD, et al. Treatment of chronic pancreatic pleural effusion by percutaneous catheter drainage of abdominal pseudocyst. *Am J Med* 1984;76:329–333.
 235. Dewan NA, Kinney WW, O'Donohue WJ Jr. Chronic massive pancreatic pleural effusion. *Chest* 1984;85: 497–501.
 236. Akahane T, Kuriyama S, Matsumoto M, et al. Pancreatic pleural effusion with a pancreaticopleural fistula diagnosed by magnetic resonance cholangiopancreatography and cured by somatostatin analogue treatment. *Abdom Imaging* 2003;28:92–95.
 237. Mori Y, Iwai A, Inagaki T, et al. Pancreaticopleural fistula imaged with magnetic resonance pancreatography. *Pancreatology* 2001;1:369–370.
 238. Materne R, Vrancx P, Pauls C, et al. Pancreaticopleural fistula: diagnosis with magnetic resonance pancreatography. *Chest* 2000;117:912–914.
 239. Glorioso LW, 3rd, Lang EK. Pulmonary manifestations of renal disease. *Radiol Clin North Am* 1984;22:647–658.
 240. Hopps H, Wissler R. Uremic pneumonitis. *Am J Pathol* 1955;31:261–274.
 241. Berger HW, Rammohan G, Neff MS, et al. Uremic pleural effusion. A study in 14 patients on chronic dialysis. *Ann Intern Med* 1975;82:362–364.
 242. Nidus BD, Matalon R, Cantacuzino D, et al. Uremic pleuritis – a clinicopathological entity. *N Engl J Med* 1969;281:255–256.
 243. Galen MA, Steinberg SM, Lowrie EG, et al. Hemorrhagic pleural effusion in patients undergoing chronic hemodialysis. *Ann Intern Med* 1975;82:359–361.
 244. Gilbert L, Ribot S, Frankel H, et al. Fibrinous uremic pleuritis: a surgical entity. *Chest* 1975;67:53–56.
 245. Rodelas R, Rakowski TA, Argy WP, et al. Fibrosing uremic pleuritis during hemodialysis. *JAMA* 1980;243:2424–2425.
 246. Salcedo JR. Urinothorax: report of 4 cases and review of the literature. *J Urol* 1986;135:805–808.
 247. Barek LB, Cigtay OS. Urinothorax – an unusual pleural effusion. *Br J Radiol* 1975;48:685–686.
 248. Baron RL, Stark DD, McClellan BL, et al. Intrathoracic extension of retroperitoneal urine collections. *AJR Am J Roentgenol* 1981;137:37–41.
 249. Lahiry SK, Alkhafaji AH, Brown AL. Urinothorax following blunt trauma to the kidney. *J Trauma* 1978;18:608–610.
 250. Redman JF, Arnold WC, Smith PL, et al. Hypertension and urinothorax following an attempted percutaneous nephrostomy. *J Urol* 1982;128: 1307–1308.
 251. Carcillo J Jr, Salcedo JR. Urinothorax as a manifestation of nondilated obstructive uropathy following renal transplantation. *Am J Kidney Dis* 1985;5:211–213.
 252. Corriere JN Jr, Miller WT, Murphy JJ. Hydronephrosis as a cause of pleural effusion. *Radiology* 1968;90:79–84.
 253. Laforet EG, Kornitzer GD. Nephrogenic pleural effusion. *J Urol* 1977;117:118–119.
 254. Nusser RA, Culhane RH. Recurrent transudative effusion with an abdominal mass. *Urinothorax*. *Chest* 1986;90: 263–264.
 255. Jenkins PG, Shelp WD. Recurrent pleural transudate in the nephrotic syndrome. A new approach to treatment. *JAMA* 1974;230:587–588.
 256. Llach F, Arieff AI, Massry SG. Renal vein thrombosis and nephrotic syndrome. A prospective study of 36 adult patients. *Ann Intern Med* 1975;83:8–14.
 257. Holzel A, Fawcitt J. Pulmonary changes in acute glomerulonephritis in childhood. *J Pediatr* 1960;57:695–703.
 258. Edwards SR, Unger AM. Acute hydrothorax – a new complication of peritoneal dialysis. *JAMA* 1967;199:853–855.
 259. Rudnick MR, Coyle JF, Beck LH, et al. Acute massive hydrothorax complicating peritoneal dialysis, report of 2 cases and a review of the literature. *Clin Nephrol* 1979;12:38–44.
 260. Finn R, Jowett EW. Acute hydrothorax complicating peritoneal dialysis. *Br Med J* 1970;2:94.
 261. Townsend R, Fragola JA. Hydrothorax in a patient receiving continuous ambulatory peritoneal dialysis: successful treatment with intermittent peritoneal dialysis. *Arch Intern Med* 1982;142:1571–1572.
 262. Nassberger L. Left-sided pleural effusion secondary to continuous ambulatory peritoneal dialysis. *Acta Med Scand* 1982;211:219–220.

263. Lorentz WB Jr. Acute hydrothorax during peritoneal dialysis. *J Pediatr* 1979;94:417-419.
264. Polsky MS, Weber CH, Ball TP Jr. Infected pyelocaliceal diverticulum and sympathetic pleural effusion. *J Urol* 1975;114:301-303.
265. Chun CH, Raff MJ, Contreras L, et al. Splenic abscess. *Medicine (Baltimore)* 1980;59:50-65.
266. Johnson JD, Raff MJ, Barnwell PA, et al. Splenic abscess complicating infectious endocarditis. *Arch Intern Med* 1983;143:906-912.
267. Sarr MG, Zuidema GD. Splenic abscess - presentation, diagnosis, and treatment. *Surgery* 1982;92:480-485.
268. Warren MS, Gibbons RB. Left-sided pleural effusion secondary to splenic vein thrombosis. A previously unrecognized relationship. *Chest* 1991;100:574-575.
269. Koehler PR, Jones R. Association of left-sided pleural effusions and splenic hematomas. *AJR Am J Roentgenol* 1980;135:851-853.
270. Light RW, George RB. Incidence and significance of pleural effusion after abdominal surgery. *Chest* 1976;69:621-625.
271. Nielsen PH, Jepsen SB, Olsen AD. Postoperative pleural effusion following upper abdominal surgery. *Chest* 1989;96:1133-1135.
272. Aronchick JM, Epstein DM, Geffer WB, et al. Chronic traumatic diaphragmatic hernia: the significance of pleural effusion. *Radiology* 1988;168:675-678.
273. Libshitz HI, Southard ME. Complications of radiation therapy: the thorax. *Semin Roentgenol* 1974;9:41-49.
274. Whitcomb ME, Schwarz MI. Pleural effusion complicating intensive mediastinal radiation therapy. *Am Rev Respir Dis* 1971;103:100-107.
275. Bachman A, Macken K. Pleural effusions following supravoltage radiation for breast carcinoma. *Radiology* 1959;72:699-709.
276. Antony VB. Drug-induced pleural disease. *Clin Chest Med* 1998;19:331-340.
277. Miller WT, Jr. Pleural and mediastinal disorders related to drug use. *Semin Roentgenol* 1995;30:35-48.
278. Urban C, Nirenberg A, Caparros B, et al. Chemical pleuritis as the cause of acute chest pain following high-dose methotrexate treatment. *Cancer* 1983;51:34-37.
279. Walden PA, Mitchell-Heggs PF, Coppin C, et al. Pleurisy and methotrexate treatment. *Br Med J* 1977;2:867.
280. Everts CS, Westcott JL, Bragg DG. Methotrexate therapy and pulmonary disease. *Radiology* 1973;107:539-543.
281. Ecker MD, Jay B, Keohane MF. Procarbazine lung. *AJR Am J Roentgenol* 1978;131:527-528.
282. Orwoll ES, Kiessling PJ, Patterson JR. Interstitial pneumonia from mitomycin. *Ann Intern Med* 1978;89:352-355.
283. Smalley RV, Wall RL. Two cases of busulfan toxicity. *Ann Intern Med* 1966;64:154-164.
284. Holoye PY, Luna MA, MacKay B, et al. Bleomycin hypersensitivity pneumonitis. *Ann Intern Med* 1978;88:47-49.
285. Pascual RS, Mosher MB, Sikand RS, et al. Effects of bleomycin on pulmonary function in man. *Am Rev Respir Dis* 1973;108:211-217.
286. Mann H, Ward JH, Samlowski WE. Vascular leak syndrome associated with interleukin-2: chest radiographic manifestations. *Radiology* 1990;176:191-194.
287. Saxon RR, Klein JS, Bar MH, et al. Pathogenesis of pulmonary edema during interleukin-2 therapy: correlation of chest radiographic and clinical findings in 54 patients. *AJR Am J Roentgenol* 1991;156:281-285.
288. Conant EF, Fox KR, Miller WT. Pulmonary edema as a complication of interleukin-2 therapy. *AJR Am J Roentgenol* 1989;152:749-752.
289. Jung JI, Choi JE, Hahn ST, et al. Radiologic features of all-trans-retinoic acid syndrome. *AJR Am J Roentgenol* 2002;178:475-480.
290. Holmberg L, Boman G. Pulmonary reactions to nitrofurantoin. 447 cases reported to the Swedish Adverse Drug Reaction Committee 1966-1976. *Eur J Respir Dis* 1981;62:180-189.
291. Pfitzenmeyer P, Foucher P, Dennewald G, et al. Pleuropulmonary changes induced by ergoline drugs. *Eur Respir J* 1996;9:1013-1019.
292. Knoop C, Mairesse M, Lenclud C, et al. Pleural effusion during bromocriptine exposure in two patients with pre-existing asbestos pleural plaques: a relationship? *Eur Respir J* 1997;10:2898-2901.
293. Graham JR, Suby HI, LeCompte PR, et al. Fibrotic disorders associated with methysergide therapy for headache. *N Engl J Med* 1966;274:359-368.
294. Graham JR. Cardiac and pulmonary fibrosis during methysergide therapy for headache. *Am J Med Sci* 1967;254:1-12.
295. Hindle W, Posner E, Sweetnam MT, et al. Pleural effusion and fibrosis during treatment with methysergide. *Br Med J* 1970;1:605-606.
296. Kok-Jensen A, Lindeneg O. Pleurisy and fibrosis of the pleura during methysergide treatment of hemigranias. *Scand J Respir Dis* 1970;51:218-222.
297. Geffer WB, Epstein DM, Bonavita JA, et al. Pleural thickening caused by Sansert and Ergotrate in the treatment of migraine. *AJR Am J Roentgenol* 1980;135:375-377.
298. Taal BG, Spierings EL, Hilvering C. Pleuropulmonary fibrosis associated with chronic and excessive intake of ergotamine. *Thorax* 1983;38:396-398.
299. Allen MB, Tosh G, Walters G, et al. Pleural and pericardial fibrosis after ergotamine therapy. *Respir Med* 1994;88:67-69.
300. Cooper JA Jr, White DA, Matthay RA. Drug-induced pulmonary disease. Part 2: Noncytotoxic drugs. *Am Rev Respir Dis* 1986;133:488-505.
301. Kuhlman J, Teigen C, Ren H, Hruban R, et al. Amiodarone pulmonary toxicity: CT findings in symptomatic patients. *Radiology* 1990;177:121-125.
302. Gonzalez-Rothi RJ, Hannan SE, Hood CI, et al. Amiodarone pulmonary toxicity presenting as bilateral exudative pleural effusions. *Chest* 1987;92:179-182.
303. Rakita L, Sobol SM, Mostow N, et al. Amiodarone pulmonary toxicity. *Am Heart J* 1983;106:906-916.
304. Zaher C, Hamer A, Peter T, et al. Low-dose steroid therapy for prophylaxis of amiodarone-induced pulmonary infiltrates. *N Engl J Med* 1983;308:779.
305. Middleton KL, Santella R, Couser JI Jr. Eosinophilic pleuritis due to propylthiouracil. *Chest* 1993;103:955-956.
306. Petusevsky ML, Faling LJ, Rocklin RE, et al. Pleuropericardial reaction to treatment with dantrolene. *JAMA* 1979;242:2772-2774.
307. Webb DB, Whale RJ. Pleuropericardial effusion associated with minoxidil administration. *Postgrad Med J* 1982;58:319-320.
308. McElvaney NG, Wilcox PG, Churg A, et al. Pleuropulmonary disease during bromocriptine treatment of Parkinson's disease. *Arch Intern Med* 1988;148:2231-2236.
309. Wiggins J, Skinner C. Bromocriptine induced pleuropulmonary fibrosis. *Thorax* 1986;41:328-330.
310. Shaunak S, Wilkins A, Pilling JB, et al. Pericardial, retroperitoneal, and pleural fibrosis induced by pergolide. *J Neurol Neurosurg Psychiatry* 1999;66:79-81.
311. Erwtaman TM, Braat MC, van Aken WG. Interstitial pulmonary fibrosis: a new side effect of practolol. *Br Med J* 1977;2:297-298.
312. Fleming HA, Hickling P. Letter: Pleural effusions after practolol. *Lancet* 1975;2:1202.
313. Hall DR, Morrison JB, Edwards FR. Pleural fibrosis after practolol therapy. *Thorax* 1978;33:822-824.
314. Marshall AJ, Eltringham WK, Barritt DW, et al. Respiratory disease associated with practolol therapy. *Lancet* 1977;2:1254-1257.
315. Wood GM, Bolton RP, Muers MF, et al. Pleurisy and pulmonary granulomas

1084 PLEURA AND PLEURAL DISORDERS

- after treatment with acebutolol. *Br Med J (Clin Res Ed)* 1982;285:936.
316. Ahmad S. Sclerosing peritonitis and propranolol. *Chest* 1981;79:361-362.
 317. Jewelewicz R, Vande Wiele RL. Acute hydrothorax as the only symptom of ovarian hyperstimulation syndrome. *Am J Obstet Gynecol* 1975;121:1121.
 318. Roden S, Juvin K, Homasson JP, et al. An uncommon etiology of isolated pleural effusion. The ovarian hyperstimulation syndrome. *Chest* 2000;118:256-258.
 319. Tansutthiwong AA, Srisombut C, Rojanasakul A. Unilateral massive pleural effusion as the only principal manifestation of severe ovarian hyperstimulation syndrome. *J Assist Reprod Genet* 2000;17:454-456.
 320. Loret de Mola JR. Pathophysiology of unilateral pleural effusions in the ovarian hyperstimulation syndrome. *Hum Reprod* 1999;14:272-273.
 321. Brown SD, Brashear RE, Schnute RB. Pleural effusion in a young woman with myxedema. *Arch Intern Med* 1983;143:1458-1460.
 322. Chetty KG. Transudative pleural effusions. *Clin Chest Med* 1985;6:49-54.
 323. Gottehrer A, Roa J, Stanford GG, et al. Hypothyroidism and pleural effusions. *Chest* 1990;98:1130-1132.
 324. Marks P, Roof B. Pericardial effusion associated with myxedema. *Ann Intern Med* 1953;39:230-240.
 325. Schneiersen S, Katz M. Solitary pleural effusion due to myxedema. *JAMA* 1958;168:1003-1005.
 326. Samman P, White W. The "yellow nail" syndrome. *Br J Dermatol* 1964;76:153-157.
 327. Hurwitz P, Pinals D. Pleural effusion in chronic hereditary lymphedema (Nonne, Milroy, Meige's disease). *Radiology* 1964;82:246-248.
 328. Emerson PA. Yellow nails, lymphoedema, and pleural effusions. *Thorax* 1966;21:247-253.
 329. Nordkild P, Kromann-Andersen H, Struve-Christensen E. Yellow nail syndrome - the triad of yellow nails, lymphedema and pleural effusions. A review of the literature and a case report. *Acta Med Scand* 1986;219:221-227.
 330. Beer DJ, Pereira W Jr, Snider GL. Pleural effusion associated with primary lymphedema: a perspective on the yellow nail syndrome. *Am Rev Respir Dis* 1978;117:595-599.
 331. Anon. Yellow nails and oedema. *Br Med J* 1972;4:130.
 332. Hiller E, Rosenow EC 3rd, Olsen AM. Pulmonary manifestations of the yellow nail syndrome. *Chest* 1972;61:452-458.
 333. Wiggins J, Strickland B, Chung KF. Detection of bronchiectasis by high-resolution computed tomography in the yellow nail syndrome. *Clin Radiol* 1991;43:377-379.
 334. Solal-Celigny P, Cormier Y, Fournier M. The yellow nail syndrome. Light and electron microscopic aspects of the pleura. *Arch Pathol Lab Med* 1983;107:183-185.
 335. Runyon BA, Forker EL, Sopko JA. Pleural-fluid kinetics in a patient with primary lymphedema, pleural effusions, and yellow nails. *Am Rev Respir Dis* 1979;119:821-825.
 336. Glazer M, Berkman N, Lafair JS, et al. Successful talc slurry pleurodesis in patients with nonmalignant pleural effusion. *Chest* 2000;117:1404-1409.
 337. Jiva TM, Poe RH, Kallay MC. Pleural effusion in yellow nail syndrome: chemical pleurodesis and its outcome. *Respiration* 1994;61:300-302.
 338. Cook GC. Periodic disease, recurrent polyserositis, familial Mediterranean fever, or simply 'FMF'. *Q J Med* 1986;60:819-823.
 339. Barakat MH, Karnik AM, Majeed HW, et al. Familial Mediterranean fever (recurrent hereditary polyserositis) in Arabs - a study of 175 patients and review of the literature. *Q J Med* 1986;60:837-847.
 340. Nugent FW, Burns JR. Periodic disease. *Med Clin North Am* 1966;50:371-378.
 341. Sohar E, Gafni J, Pras M, et al. Familial Mediterranean fever. A survey of 470 cases and review of the literature. *Am J Med* 1967;43:227-253.
 342. Mancini JL. Familial paroxysmal polyserositis, phenotype I (familial Mediterranean fever). A rare cause of pleurisy. Case report and review of the literature. *Am Rev Respir Dis* 1973;107:461-463.
 343. el-Kassimi FA. Acute pleuritic chest pain with pleural effusion and plate atelectasis. Familial Mediterranean fever (periodic disease). *Chest* 1987;91:265-266.
 344. Meyerhoff J. Familial Mediterranean fever: report of a large family, review of the literature, and discussion of the frequency of amyloidosis. *Medicine (Baltimore)* 1980;59:66-77.
 345. Villena V, Lopez-Encuentra A, Pozo F, et al. Measurement of pleural pressure during therapeutic thoracentesis. *Am J Respir Crit Care Med* 2000;162:1534-1538.
 346. Light RW, Jenkinson SG, Minh VD, et al. Observations on pleural fluid pressures as fluid is withdrawn during thoracentesis. *Am Rev Respir Dis* 1980;121:799-804.
 347. Bessone LN, Ferguson TB, Burford TH. Chylothorax. *Ann Thorac Surg* 1971;12:527-550.
 348. Dahlgren S. Anatomy of the thoracic duct from the standpoint of surgery for chylothorax. *Acta Chir Scand* 1963;125:201-206.
 349. Rosenberger A, Abrams HL. Radiology of the thoracic duct. *Am J Roentgenol Radium Ther Nucl Med* 1971;111:807-820.
 350. Sassoon CS, Light RW. Chylothorax and pseudochylothorax. *Clin Chest Med* 1985;6:163-171.
 351. Schulman A, Fataar S, Dalrymple R, et al. The lymphographic anatomy of chylothorax. *Br J Radiol* 1978;51:420-427.
 352. Kausel H, Reeve T, Stein A, et al. Anatomic and pathologic studies of the thoracic duct. *J Thorac Surg* 1957;34:631-641.
 353. Van Pernis P. Variations of thoracic duct. *Surgery* 1949;26:806-809.
 354. Meade R, Head J, CW M. The management of chylothorax. *J Thorac Surg* 1950;19:709-723.
 355. Thorne P. Traumatic chylothorax. *Tubercle* 1958;39:29-34.
 356. Neyazaki T, Kupic EA, Marshall WH, et al. Collateral lymphatico-venous communications after experimental obstruction of the thoracic duct. *Radiology* 1965;85:423-432.
 357. Lampson R. Traumatic chylothorax: a review of the literature and report of a case treated by mediastinal ligation of the thoracic duct. *J Thorac Surg* 1948;17:778-791.
 358. Meade R, Head J, Moen C. The management of chylothorax. *J Thorac Surg* 1950;19:709-723.
 359. Valentine VG, Raffin TA. The management of chylothorax. *Chest* 1992;102:586-591.
 360. Grant T, Levin B. Lymphangiographic visualization of pleural and pulmonary lymphatics in a patient without chylothorax. *Radiology* 1974;113:49-50.
 361. Trapnell D. The peripheral lymphatics of the lung. *Br J Radiol* 1963;36:660-672.
 362. Weidner WA, Steiner RM. Roentgenographic demonstration of intrapulmonary and pleural lymphatics during lymphangiography. *Radiology* 1971;100:533-539.
 363. Ducharme JC, Belanger R, Simard P, et al. Chylothorax, chylopericardium with multiple lymphangioma of bone. *J Pediatr Surg* 1982;17:365-367.
 364. Berkenbosch JW, Monteleone PM, Tobias JD. Chylothorax following apparently spontaneous central venous thrombosis in a patient with septic shock. *Pediatr Pulmonol* 2003;35:230-233.
 365. Kramer SS, Taylor GA, Garfinkel DJ, et al. Lethal chylothoraces due to superior vena caval thrombosis in infants. *AJR Am J Roentgenol* 1981;137:559-563.
 366. Thurer RJ. Chylothorax: a complication of subclavian vein catheterization and parenteral hyperalimentation. *J Thorac Cardiovasc Surg* 1976;71:465-468.
 367. Fossum TW, Birchard SJ. Lymphangiographic evaluation of experimentally induced chylothorax

- after ligation of the cranial vena cava in dogs. *Am J Vet Res* 1986;47:967-971.
368. Nix J, Albert M, Dugas J, et al. Chylothorax and chylous ascites; a study of 302 selected cases. *Am J Gastroenterol* 1957;28:40-53.
 369. Ross J. A review of the surgery of the thoracic duct. *Thorax* 1961;16:12-21.
 370. Bower G. Chylothorax: observations in 20 cases. *Dis Chest* 1964;46:464-468.
 371. Staats BA, Ellefson RD, Budahn LL, et al. The lipoprotein profile of chylous and nonchylous pleural effusions. *Mayo Clin Proc* 1980;55:700-704.
 372. Seriff NS, Cohen ML, Samuel P, et al. Chylothorax: diagnosis by lipoprotein electrophoresis of serum and pleural fluid. *Thorax* 1977;32:98-100.
 373. Yeam I, Sassoon C. Hemothorax and chylothorax. *Curr Opin Pulm Med* 1997;3:310-314.
 374. Macfarlane JR, Holman CW. Chylothorax. *Am Rev Respir Dis* 1972;105:287-291.
 375. Roy P, Carr D, Payne W. The problem of chylothorax. *Mayo Clin Proc* 1967;42:457-467.
 376. Cevese PG, Vecchioni R, D'Amico DF, et al. Postoperative chylothorax. Six cases in 2,500 operations, with a survey of the world literature. *J Thorac Cardiovasc Surg* 1975;69:966-971.
 377. Kshetry VR, Rebello R. Chylothorax after coronary artery bypass grafting. *Thorax* 1982;37:954.
 378. Weber DO, Mastro PD, Yarnoz MD. Chylothorax after myocardial revascularization with internal mammary graft. *Ann Thorac Surg* 1981;32:499-492.
 379. Nygaard SD, Berger HA, Fick RB. Chylothorax as a complication of oesophageal sclerotherapy. *Thorax* 1992;47:134-135.
 380. Shimizu K, Yoshida J, Nishimura M, et al. Treatment strategy for chylothorax after pulmonary resection and lymph node dissection for lung cancer. *J Thorac Cardiovasc Surg* 2002;124:499-502.
 381. Orringer MB, Bluett M, Deeb GM. Aggressive treatment of chylothorax complicating transhiatal esophagectomy without thoracotomy. *Surgery* 1988;104:720-726.
 382. Dulchavsky SA, Ledgerwood AM, Lucas CE. Management of chylothorax after blunt chest trauma. *J Trauma* 1988;28:1400-1401.
 383. Cammarata SK, Brush RE Jr, Hyzy RC. Chylothorax after childbirth. *Chest* 1991;99:1539-1540.
 384. Reilly KM, Tsou E. Bilateral chylothorax. A case report following episodes of stretching. *JAMA* 1975;233:536-537.
 385. van Straaten HL, Gerards LJ, Krediet TG. Chylothorax in the neonatal period. *Eur J Pediatr* 1993;152:2-5.
 386. Ryu JH, Doerr CH, Fisher SD, et al. Chylothorax in lymphangi-oleiomyomatosis. *Chest* 2003;123:623-627.
 387. Gardner TW, Domm AC, Brock CE, et al. Congenital pulmonary lymphangiectasis. A case complicated by chylothorax. *Clin Pediatr (Phila)* 1983;22:75-78.
 388. Baltaxe HA, Lee JG, Ehlers KH, et al. Pulmonary lymphangiectasia demonstrated by lymphangiography in 2 patients with Noonan's syndrome. *Radiology* 1975;115:149-153.
 389. Brown LR, Reiman HM, Rosenow EC 3rd, et al. Intrathoracic lymphangioma. *Mayo Clin Proc* 1986;61:882-892.
 390. Duckett JG, Lazarus A, White KM. Cutaneous masses, rib lesions, and chylous pleural effusion in a 20-year-old man. *Chest* 1990;97:1227-1228.
 391. Steiner GM, Farman J, Lawson JP. Lymphangiomatosis of bone. *Radiology* 1969;93:1093-1098.
 392. Aviv R, McHugh K. Mechanisms of chylous effusion in lymphangiomatosis. *AJR Am J Roentgenol* 2000;175:1191.
 393. Swensen SJ, Hartman TE, Mayo JR, et al. Diffuse pulmonary lymphangiomatosis: CT findings. *J Comput Assist Tomog* 1995;19:348-352.
 394. Takahashi K, Takahashi H, Maeda K, et al. An adult case of lymphangiomatosis of the mediastinum, pulmonary interstitium and retroperitoneum complicated by chronic disseminated intravascular coagulation. *Eur Respir J* 1995;8:1799-1802.
 395. Lee WS, Kim SH, Kim I, et al. Chylothorax in Gorham's disease. *J Korean Med Sci* 2002;17:826-829.
 396. Yoo SY, Goo JM, Im JG. Mediastinal lymphangioma and chylothorax: thoracic involvement of Gorham's disease. *Korean J Radiol* 2002;3:130-132.
 397. Brenner WI, Boal BH, Reed GE. Chylothorax as a manifestation of rheumatic mitral stenosis: its postoperative management with a diet of medium-chain triglycerides. *Chest* 1978;73:672-673.
 398. Villena V, de Pablo A, Martin-Escribano P. Chylothorax and chylous ascites due to heart failure. *Eur Respir J* 1995;8:1235-1236.
 399. Effmann EL, Ablow RC, Touloukian RJ, et al. Radiographic aspects of total parenteral nutrition during infancy. *Radiology* 1978;127:195-201.
 400. Hinckley ME. Thoracic-duct thrombosis with fatal chylothorax caused by a long venous catheter. *N Engl J Med* 1969;280:95-96.
 401. Warren WH, Altman JS, Gregory SA. Chylothorax secondary to obstruction of the superior vena cava: a complication of the LeVeen shunt. *Thorax* 1990;45:978-979.
 402. Adams J, McEvoy R, DeWeese J. Primary deep venous thrombosis of upper extremity. *Arch Surg* 1965;91:29-42.
 403. Delgado C, Martin M, de la Portilla F. Retrosternal goiter associated with chylothorax. *Chest* 1994;106:1924-1925.
 404. Vennera MC, Moreno R, Cot J, et al. Chylothorax and tuberculosis. *Thorax* 1983;38:694-695.
 405. Freundlich IM. The role of lymphangiography in chylothorax. A report of six nontraumatic cases. *Am J Roentgenol Radium Ther Nucl Med* 1975;125:617-627.
 406. Kitchen ND, Hocken DB, Greenhalgh RM, et al. Use of the Denver pleuroperitoneal shunt in the treatment of chylothorax secondary to filariasis. *Thorax* 1991;46:144-145.
 407. Moss R, Hinds S, Fedullo AJ. Chylothorax: a complication of the nephrotic syndrome. *Am Rev Respir Dis* 1989;140:1436-1437.
 408. Valdes L, Alvarez D, Pose A, et al. Cirrhosis of the liver, an exceptional cause of chylothorax: two cases. *Respir Med* 1996;90:61-62.
 409. Romero S, Martin C, Hernandez L, et al. Chylothorax in cirrhosis of the liver: analysis of its frequency and clinical characteristics. *Chest* 1998;114:154-159.
 410. Jarman PR, Whyte MK, Sabroe I, et al. Sarcoidosis presenting with chylothorax. *Thorax* 1995;50:1324-1325.
 411. Riantawan P, Tansupasawasdikul S, Subhannachart P. Bilateral chylothorax complicating massive osteolysis (Gorham's syndrome). *Thorax* 1996;51:1277-1278.
 412. Anton PA, Rubio J, Casan P, et al. Chylothorax due to *Mycobacterium tuberculosis*. *Thorax* 1995;50:1019.
 413. Lowell J. Pleural effusions: a comprehensive review. Baltimore: University Park Press, 1977.
 414. Fairfax AJ, McNabb WR, Spiro SG. Chylothorax: a review of 18 cases. *Thorax* 1986;41:880-885.
 415. Hom M, Jolles H. Traumatic mediastinal lymphocoele mimicking other thoracic injuries: case report. *J Thorac Imaging* 1992;7:78-80.
 416. Suzuki K, Yoshida J, Nishimura M, et al. Postoperative mediastinal chyloma. *Ann Thorac Surg* 1999;68:1857-1858.
 417. Higgins CB, Mulder DG. Mediastinal chyloma, a roentgenographic sign of chylous fistula. *JAMA* 1970;211:1188.
 418. Mine H, Tamura K, Tanegashima K, et al. Non-traumatic chylothorax associated with diffuse lymphatic dysplasia. *Lymphology* 1984;17:111-112.
 419. Ngan H, Fok M, Wong J. The role of lymphography in chylothorax following thoracic surgery. *Br J Radiol* 1988;61:1032-1036.
 420. Day DL, Warwick WJ. Thoracic duct opacification for CT scanning. *AJR Am J Roentgenol* 1985;144:403-404.
 421. Qureshy A, Kubota K, Ono S, et al. Thoracic duct scintigraphy by orally

1086 PLEURA AND PLEURAL DISORDERS

- administered I-123 BMIPP: normal findings and a case report. *Clin Nucl Med* 2001;26:847-855.
422. Kettner BI, Aurisch R, Ruckert JC, et al. Scintigraphic localization of lymphatic leakage site after oral administration of iodine-123-IPPA. *J Nucl Med* 1998;39:2141-2144.
 423. Al-Sebeih K, Sadeghi N, Al-Dahri S. Bilateral chylothorax following neck dissection: a new method of treatment. *Ann Otol Rhinol Laryngol* 2001;110:381-384.
 424. Hillerdal G. Chylothorax and pseudochylothorax. *Eur Respir J* 1997;10:1157-1162.
 425. Hillerdal G. Chyliform (cholesterol) pleural effusion. *Chest* 1985;88:426-428.
 426. Ferguson GC. Cholesterol pleural effusion in rheumatoid lung disease. *Thorax* 1966;21:577-582.
 427. Johnson JR, Falk A, Iber C, et al. Paragonimiasis in the United States. A report of nine cases in Hmong immigrants. *Chest* 1982;82:168-171.
 428. Groskin SA. Selected topics in chest trauma. *Radiology* 1992;183:605-617.
 429. Banks J, Cassidy D, Campbell IA, et al. Unusual clinical signs complicating tension haemothorax. *Br J Dis Chest* 1984;78:272-274.
 430. Willson SA, Sawicka EH, Mitchell IC. Spontaneous pneumothorax: an unusual radiological appearance. *Br J Radiol* 1985;58:173-175.
 431. Milner LB, Ryan K, Gullo J. Fatal intrathoracic hemorrhage after percutaneous aspiration lung biopsy. *AJR* 1979;132:280-281.
 432. Rodriguez E, Martinez J, Javaloyas M, et al. Haemothorax in the course of chickenpox. *Thorax* 1986;41:491.
 433. Rasaretnam R, Chanmugam D, Sivathanan C. Spontaneous haemothorax in a mild haemophilic. *Thorax* 1976;31:601-604.
 434. Millard CE. Massive haemothorax complicating heparin therapy for pulmonary infarction. *Chest* 1971;59:235-237.
 435. Robinson NMK, Thomas MR, Jewitt DE. Spontaneous haemothorax as a complication of anti-coagulation following coronary angioplasty. *Respir Med* 1995;89:629-630.
 436. Spear BS, Sully L, Lewis CT. Pulmonary arteriovenous fistula presenting as spontaneous haemothorax. *Thorax* 1975;30:355-356.
 437. DeFrance JH, Blewett JH, Ricci JA, et al. Massive haemothorax: two unusual cases. *Chest* 1974;66:82-84.
 438. Castells L, Comas P, Gonzalez A, et al. Case report: haemothorax in hereditary multiple exostosis. *Br J Radiol* 1993;66:269-270.
 439. Kravis MMJ, Hutton LC. Solitary plasma cell tumor of the pleura manifested as massive haemothorax. *AJR* 1993;161:543-544.
 440. Kupferschmid JP, Shahian DM, Villanueva AG. Massive haemothorax associated with intrathoracic extramedullary hematopoiesis involving the pleura. *Chest* 1993;103:974-975.
 441. Sulis E, Floris C. Haemothorax due to thoracic extramedullary erythropoiesis in thalassaemia intermedia. *Br Med J* 1985;291:1094.
 442. Calvert RJ, Smith E. An analytical review of spontaneous haemopneumothorax. *Thorax* 1955;10:64-72.
 443. Shepard MK, Mancini MC, Campbell GD Jr, George R. Right-sided haemothorax and recurrent abdominal pain in a 34-year-old woman. *Chest* 1993;103:1239-1240.
 444. Yeh T. Endometriosis within the thorax: metaplasia, implantation, or metastasis? *J Cardiovasc Surg* 1967;53:201-205.
 445. Deaton W, Johnston F. Spontaneous hemopneumothorax. *J Thorac Cardiovasc Surg* 1962;43:413-415.
 446. Slind RO, Rodarte JR. Spontaneous haemothorax in an otherwise healthy young man. *Chest* 1974;66:81.
 447. Yung CM, Bessen SC, Hingorani V, et al. Idiopathic haemothorax. *Chest* 1979;103:638-639.
 448. Reynolds J, Davis JT. Injuries of the chest wall, pleura, pericardium, lungs, bronchi and esophagus. *Radiol Clin North Am* 1966;4:383-401.
 449. Qureshi MM, Roble DC, Gindin RA, et al. Subarachnoid-pleural fistula. Case report and review of the literature. *J Thorac Cardiovasc Surg* 1986;91:238-241.
 450. Wolverson MK, Crepps LF, Sundaram M, et al. Hyperdensity of recent hemorrhage at body computed tomography: incidence and morphologic variation. *Radiology* 1983;148:779-784.
 451. McLoud TC, Isler R, Head J. The radiologic appearance of chemical pleurodesis. *Radiology* 1980;135:313-317.
 452. Eyler WR, Monsein LH, Beute GH, et al. Rib enlargement in patients with chronic pleural disease. *AJR Am J Roentgenol* 1996;167:921-926.
 453. Aberle DR, Gamsu G, Ray CS. High-resolution CT of benign asbestos-related diseases: clinical and radiographic correlation. *AJR Am J Roentgenol* 1988;151:883-891.
 454. Leung AN, Müller NL, Miller RR. CT in differential diagnosis of diffuse pleural disease. *AJR Am J Roentgenol* 1990;154:487-492.
 455. Im J-G, Webb W, Rosen A, et al. Costal pleura: Appearances at high-resolution CT. *Radiology* 1989;171:125-131.
 456. Hulnick DH, Naidich DP, McCauley DI. Pleural tuberculosis evaluated by computed tomography. *Radiology* 1983;149:759-765.
 457. Schmitt WG, Hubener KH, Rucker HC. Pleural calcification with persistent effusion. *Radiology* 1983;149:633-638.
 458. Waite RJ, Carbonneau RJ, Balikian JP, et al. Parietal pleural changes in empyema: appearances at CT. *Radiology* 1990;175:145-150.
 459. Takasugi JE, Godwin JD, Teefey SA. The extrapleural fat in empyema: CT appearance. *Br J Radiol* 1991;64:580-583.
 460. Bury T, Paulus P, Dowlati A, et al. Evaluation of pleural diseases with FDG-PET imaging: preliminary report. *Thorax* 1997;52:187-189.
 461. Hierholzer J, Luo L, Bittner RC, et al. MRI and CT in the differential diagnosis of pleural disease. *Chest* 2000;118:604-609.
 462. Franquet T, Gimenez A, Cremades R, et al. Spontaneous reversibility of "pleural thickening" in a patient with semi-invasive pulmonary aspergillosis: radiographic and CT findings. *Eur Radiol* 2000;10:722-724.
 463. Chong VF, Fan YF. Invasive thymoma presenting as pleural thickening. *AJR Am J Roentgenol* 1997;168:568-569.
 464. Lee MJ, Breatnach E. Pleural thickening caused by leukemic infiltration: pleural findings. *AJR Am J Roentgenol* 1994;163:1527-1528.
 465. Neff CC, vanSonnenberg E, Lawson DW, et al. CT follow-up of empyemas: pleural peels resolve after percutaneous catheter drainage. *Radiology* 1990;176:195-197.
 466. Buchanan DR, Johnston ID, Kerr IH, et al. Cryptogenic bilateral fibrosing pleuritis. *Br J Dis Chest* 1988;82:186-193.
 467. Lee-Chiong TL Jr, Hilbert J. Extensive idiopathic benign bilateral asynchronous pleural fibrosis. *Chest* 1996;109:564-565.
 468. Oliver RM, Neville E. Progressive apical pleural fibrosis: a 'constrictive' ventilatory defect. *Br J Dis Chest* 1988;82:439-443.
 469. Azoulay E, Paugam B, Heymann MF, et al. Familial extensive idiopathic bilateral pleural fibrosis. *Eur Respir J* 1999;14:971-973.
 470. Hayes JP, Wiggins J, Ward K, et al. Familial cryptogenic fibrosing pleuritis with Fanconi's syndrome (renal tubular acidosis). A new syndrome. *Chest* 1995;107:576-578.
 471. Gilmartin D. The serratus anterior muscle on chest radiographs. *Radiology* 1979;131:629-635.
 472. Collins JD, Brown RK, Batra P. Asbestosis and the serratus anterior muscle. *J Natl Med Assoc* 1983;75:296-300.
 473. Sargent EN, Boswell WD Jr, Ralls PW, et al. Subpleural fat pads in patients exposed to asbestos: distinction from non-calcified pleural plaques. *Radiology* 1984;152:273-277.
 474. Vix VA. Extrapleural costal fat. *Radiology* 1974;112:563-565.

75. Renner RR, Pernice NJ. The apical cap. *Semin Roentgenol* 1977;12:299-302.
76. Renner RR, Markarian B, Pernice NJ, et al. The apical cap. *Radiology* 1974;110:569-573.
71. Jamison H. Anatomic-roentgenographic study of pleural domes and pulmonary apices, with special reference to apical subpleural scars. *Radiology* 1941;36:302-314.
78. Butler C, 2nd, Kleinerman J. The pulmonary apical cap. *Am J Pathol* 1970;60:205-216.
79. Yousem SA. Pulmonary apical cap: a distinctive but poorly recognized lesion in pulmonary surgical pathology. *Am J Surg Pathol* 2001;25:679-683.
80. McLoud TC, Isler RJ, Novelline RA, et al. The apical cap. *AJR Am J Roentgenol* 1981;137:299-306.
81. Proto AV. Conventional chest radiographs: anatomic understanding of newer observations. *Radiology* 1992;183:593-603.
82. Im JG, Webb WR, Han MC, et al. Apical opacity associated with pulmonary tuberculosis: high-resolution CT findings. *Radiology* 1991;178:727-731.
83. Reeder LB. Malignant pleural effusions. *Curr Treat Options Oncol* 2001;2:93-96.
84. Sahn SA. Management of malignant pleural effusions. *Monaldi Arch Chest Dis* 2001;56:394-399.
85. de Campos JR, Vargas FS, de Campos Werebe E, et al. Thoracoscopy talc poudrage: a 15-year experience. *Chest* 2001;119:801-806.
86. Tschopp JM, Frey JG. Treatment of primary spontaneous pneumothorax by simple talcage under medical thoracoscopy. *Monaldi Arch Chest Dis* 2002;57:88-92.
87. Marom EM, Erasmus JJ, Herndon JE, 2nd, et al. Usefulness of imaging-guided catheter drainage and talc sclerotherapy in patients with metastatic gynecologic malignancies and symptomatic pleural effusions. *AJR Am J Roentgenol* 2002;179:105-108.
88. Marom EM, Patz EF Jr, Erasmus JJ, et al. Malignant pleural effusions: treatment with small-bore-catheter thoracostomy and talc pleurodesis. *Radiology* 1999;210:277-281.
89. Murray JG, Patz EF Jr, Erasmus JJ, et al. CT appearance of the pleural space after talc pleurodesis. *AJR Am J Roentgenol* 1997;169:89-91.
90. Light RW. Talc should not be used for pleurodesis. *Am J Respir Crit Care Med* 2000;162:2024-2026.
91. Sahn SA. Talc should be used for pleurodesis. *Am J Respir Crit Care Med* 2000;162:2023-2024; discussion 2026.
92. Brant A, Eaton T. Serious complications with talc slurry pleurodesis. *Respirology* 2001;6:181-185.
493. Vix VA. Roentgenographic manifestations of pleural disease. *Semin Roentgenol* 1977;12:277-286.
494. Sargent EN, Jacobson G, Gordonson JS. Pleural plaques: a signpost of asbestos dust inhalation. *Semin Roentgenol* 1977;12:287-297.
495. Shapir J, Lisbona A, Palayew MJ. Chronic calcified empyema. *J Can Assoc Radiol* 1981;32:24-27.
496. Im JG, Chung JW, Han MC. Milk of calcium pleural collections: CT findings. *J Comput Assist Tomogr* 1993;17:613-616.
497. Scales FE, Lee ME. Nonoperative diagnosis of intrathoracic splenosis. *AJR Am J Roentgenol* 1983;141:1273-1274.
498. Dalton ML Jr, Strange WH, Downs EA. Intrathoracic splenosis. Case report and review of the literature. *Am Rev Respir Dis* 1971;103:827-830.
499. Normand JP, Rioux M, Dumont M, et al. Thoracic splenosis after blunt trauma: frequency and imaging findings. *AJR Am J Roentgenol* 1993;161:739-741.
500. Moncada R, Williams V, Fareed J, et al. Thoracic splenosis. *AJR Am J Roentgenol* 1985;144:705-706.
501. Yamine JN, Yatim A, Barbari A. Radionuclide imaging in thoracic splenosis and a review of the literature. *Clin Nucl Med* 2003;28:121-123.
502. Naylor MF, Karstaedt N, Finck SJ, et al. Noninvasive methods of diagnosing thoracic splenosis. *Ann Thorac Surg* 1999;68:243-244.
503. Davis HH, 2nd, Varki A, Heaton WA, et al. Detection of accessory spleens with indium 111-labeled autologous platelets. *Am J Hematol* 1980;8:81-86.
504. Chee CB, Abisheganaden J, Yeo JK, et al. Persistent air-leak in spontaneous pneumothorax - clinical course and outcome. *Respir Med* 1998;92:757-761.
505. Abolnik IZ, Lossos IS, Gillis D, et al. Primary spontaneous pneumothorax in men. *Am J Med Sci* 1993;305:297-303.
506. Killen D, Gobbel W. Spontaneous pneumothorax. Boston: Little, Brown, 1968.
507. Inouye WY, Berggren RB, Johnson J. Spontaneous pneumothorax: treatment and mortality. *Dis Chest* 1967;51:67-73.
508. Melton LJ, 3rd, Hepper NG, Offord KP. Incidence of spontaneous pneumothorax in Olmsted County, Minnesota: 1950 to 1974. *Am Rev Respir Dis* 1979;120:1379-1382.
509. Greene R, McLoud TC, Stark P. Pneumothorax. *Semin Roentgenol* 1977;12:313-325.
510. Ruckley CV, McCormack RJ. The management of spontaneous pneumothorax. *Thorax* 1966;21:139-144.
511. Lesur O, Delorme N, Fromaget JM, et al. Computed tomography in the etiologic assessment of idiopathic spontaneous pneumothorax. *Chest* 1990;98:341-347.
512. Bense L, Lewander R, Eklund G, et al. Nonsmoking, non-alpha 1-antitrypsin deficiency-induced emphysema in nonsmokers with healed spontaneous pneumothorax, identified by computed tomography of the lungs [see comments]. *Chest* 1993;103:433-438.
513. Warner BW, Bailey WW, Shipley RT. Value of computed tomography of the lung in the management of primary spontaneous pneumothorax. *Am J Surg* 1991;162:39-42.
514. West JB. Distribution of mechanical stress in the lung, a possible factor in localisation of pulmonary disease. *Lancet* 1971;1:839-841.
515. Glazier JB, Hughes JM, Maloney JE, et al. Vertical gradient of alveolar size in lungs of dogs frozen intact. *J Appl Physiol* 1967;23:694-705.
516. Melton LJ, 3rd, Hepper NG, Offord KP. Influence of height on the risk of spontaneous pneumothorax. *Mayo Clin Proc* 1981;56:678-682.
517. Giuffre B. Supine pneumothoraces in adults. *Australas Radiol* 1984;28:335-338.
518. Scott GC, Berger R, McKean HE. The role of atmospheric pressure variation in the development of spontaneous pneumothoraces. *Am Rev Respir Dis* 1989;139:659-662.
519. Gilmartin JJ, Wright AJ, Gibson GJ. Effects of pneumothorax or pleural effusion on pulmonary function. *Thorax* 1985;40:60-65.
520. Norris RM, Jones JG, Bishop JM. Respiratory gas exchange in patients with spontaneous pneumothorax. *Thorax* 1968;23:427-433.
521. Kircher L, Swartzel R. Spontaneous pneumothorax and its treatment. *JAMA* 1954;155:24-29.
522. Northfield TC. Oxygen therapy for spontaneous pneumothorax. *Br Med J* 1971;4:86-88.
523. Lippert HL, Lund O, Blegvad S, et al. Independent risk factors for cumulative recurrence rate after first spontaneous pneumothorax. *Eur Respir J* 1991;4:324-331.
524. Jenkinson SG. Pneumothorax. *Clin Chest Med* 1985;6:153-161.
525. Sharpe IK, Ahmad M, Braun W. Familial spontaneous pneumothorax and HLA antigens. *Chest* 1980;78:264-268.
526. Leites V, Tannenbaum E. Familial spontaneous pneumothorax. *Am Rev Respir Dis* 1960;82:240-241.
527. Wilson WG, Aylsworth AS. Familial spontaneous pneumothorax. *Pediatrics* 1979;64:172-175.
528. Sugiyama Y, Maeda H, Yotsumoto H, et al. Familial spontaneous pneumothorax. *Thorax* 1986;41:969-970.
529. Rashid A, Sendi A, Al-Kadhimi A, et al. Concurrent spontaneous pneumothorax in identical twins. *Thorax* 1986;41:971.